


Patient Name : Ms. RIZWANA	Visit No : CHA250034675
Age/Gender : 27 Y 1 M 4 D/F	Registration ON : 26/Feb/2025 11:45AM
Lab No : 10131971	Sample Collected ON : 26/Feb/2025 11:49AM
Referred By : Dr. RUKSHANA KHAN, GREEN CROS	Sample Received ON : 26/Feb/2025 12:08PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 26/Feb/2025 01:45PM
Doctor Advice : C SPINE AP /LAT, URIC ACID, RF FACTOR, CRP (Quantitative)	



Test Name	Result	Unit	Bio. Ref. Range	Method
RF FACTOR				
RHEUMATOID FACTOR	9.70	IU/ml	0 - 14	

SUMMARY : Rheumatoid factors (RF) group of autoantibodies belonging to all immunoglobulin classes directed against the FC fragment of altered or complexed Igg. Diagnostic test for RF determination identify mainly RF of the IgM class which are detectable in several rheumatic diseases, mainly of inflammatory origin.

RF occur in approx 70 -80 % of patients with rheumatoid arthritis (RA), but they are not specific for RA as elevated concentrations are also observed in various non rheumatic disease & in approx 10 % of the elderly population without clinical symptoms of RA. High RF concentrations in RA are often associated with a more progressive clinical course of the disease. However, a positive RF value has to be confirmed by clinical & other laboratory findings.



CHARAK

[Checked By]

Print.Date/Time: 26-02-2025 15:30:51

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Ms. RIZWANA	Visit No : CHA250034675
Age/Gender : 27 Y 1 M 4 D/F	Registration ON : 26/Feb/2025 11:45AM
Lab No : 10131971	Sample Collected ON : 26/Feb/2025 11:49AM
Referred By : Dr. RUKSHANA KHAN, GREEN CROS	Sample Received ON : 26/Feb/2025 12:08PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 26/Feb/2025 12:59PM
Doctor Advice : C SPINE AP /LAT, URIC ACID, RF FACTOR, CRP (Quantitative)	



Test Name	Result	Unit	Bio. Ref. Range	Method
CRP-QUANTITATIVE				

CRP-QUANTITATIVE TEST 0.167 MG/L 0.1 - 6

Method: Immunoturbidimetric

(Method: Immunoturbidimetric on photometry system)

SUMMARY : C - reactive protien (CRP) is the best known among the acute phase protiens, a group of protien whose concentration increases in blood as a response to inflammatory disorders. CRP is normally present in low concentration in blood of healthy individuals (< 1mg/L). It is elevated up to 500 mg/L in acute inflammatory processes associated with bacterial infections, post operative conditions tissue damage already after 6 hours reaching a peak at 48 hours. The measurment of CRP represents a useful aboratory test for detection of acute infection as well as for monitoring inflammtory proceses also in acute rheumatic & gastrointestinal disease. In recent studies it has been shows that in apparrently healthy subjects there is a direct orrelation between CRP concentrations & the risk of developing oronary heart disease (CHD).

hsCRP cut off for risk assessment as per CDC/AHA Level	Risk
<1.0	Low
1.0-3.0	Average
>3.0	High

All reports to be clinically corelated

URIC ACID				
Sample Type : SERUM				
SERUM URIC ACID	4.1	mg/dL	2.40 - 5.70	Uricase, Colorimetric

*** End Of Report ***

[Checked By]

Print.Date/Time: 26-02-2025 15:30:54

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name	: Ms.RIZWANA	Visit No	: CHA250034675
Age/Gender	: 27 Y 1 M 4 D/F	Registration ON	: 26/Feb/2025 11:37AM
Lab No	: 10131971	Sample Collected ON	: 26/Feb/2025 11:37AM
Referred By	: Dr.RUKSHANA KHAN, GREEN CROSS	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 26/Feb/2025 02:49PM

SKIAGRAM CERVICAL SPINE AP AND LATERAL VIEW

- Shape and size of bodies of cervical vertebrae are normal.
- Intervertebral disc spaces are maintained.
- Posterior elements are seen normally.
- No cervical rib is seen.
- No CVJ anomaly is seen.

Clinical correlation is necessary.

[DR. RAJESH KUMAR SHARMA, MD]

TRANSCRIBED BY: ANUP

*** End Of Report ***

