

Patient Name : Ms.SHAHZADA	Visit No : CHA250035108
Age/Gender : 51 Y/F	Registration ON : 27/Feb/2025 07:07AM
Lab No : 10132404	Sample Collected ON : 27/Feb/2025 07:08AM
Referred By : Dr.U1	Sample Received ON : 27/Feb/2025 07:59AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 27/Feb/2025 12:11PM
Doctor Advice : PP,FASTING,25 OH vit. D,HBA1C (EDTA),RF FACTOR	



Test Name	Result	Unit	Bio. Ref. Range	Method
HBA1C				
Glycosylated Hemoglobin (HbA1c)	8.4	%	4 - 5.7	HPLC (EDTA)

NOTE:-

Glycosylated Hemoglobin Test (HbA1c) is performed in this laboratory by the Gold Standard Reference method, ie: HPLC Technology (High performance Liquid Chromatography D10) from Bio-Rad Laboratories. USA.

EXPECTED (RESULT) RANGE :

Bio system	Degree of normal
4.0 - 5.7 %	Normal Value (OR) Non Diabetic
5.8 - 6.4 %	Pre Diabetic Stage
> 6.5 %	Diabetic (or) Diabetic stage
6.5 - 7.0 %	Well Controlled Diabet
7.1 - 8.0 %	Unsatisfactory Control
> 8.0 %	Poor Control and needs treatment

CHARAK

[Checked By]



Print.Date/Time: 27-02-2025 12:51:51

*Patient Identity Has Not Been Verified. Not For Medicolegal

Sharma

DR. NISHANT SHARMA PATHOLOGIST
DR. SHADAB PATHOLOGIST
Dr. SYED SAIF AHMAD MD (MICROBIOLOGY)


Patient Name : Ms.SHAHZADA	Visit No : CHA250035108
Age/Gender : 51 Y/F	Registration ON : 27/Feb/2025 07:07AM
Lab No : 10132404	Sample Collected ON : 27/Feb/2025 07:08AM
Referred By : Dr.U1	Sample Received ON : 27/Feb/2025 07:59AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 27/Feb/2025 10:56AM
Doctor Advice : PP,FASTING,25 OH vit. D,HBA1C (EDTA),RF FACTOR	



Test Name	Result	Unit	Bio. Ref. Range	Method
RF FACTOR				
RHEUMATOID FACTOR	5.60	IU/ml	0 - 14	

SUMMARY : Rheumatoid factors (RF) group of autoantibodies belonging to all immunoglobulin classes directed against the FC fragment of altered or complexed Igg .Diagnostic test for RF determination identify mainly RF of the IgM class which are detectable in several rheumatic diseases, mainly of inflammatory origin.

RF occur in approx 70 -80 % of patients with rheumatoid arthritis (RA), but they are not specific for RA as elevated concentrations are also observed in various non rheumatic disease & in approx 10 % of the elderly population without clinical symptoms of RA. High RF concentrations in RA are often associated with a more progressive clinical course of the disease .However,a positive RF value has to be confirmed by clinical & other laboratory findings.



CHARAK

[Checked By]

Print.Date/Time: 27-02-2025 12:51:53

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST


Dr. SYED SAIF AHMAD
MD (MICROBIOLOGY)

Patient Name : Ms.SHAHZADA	Visit No : CHA250035108
Age/Gender : 51 Y/F	Registration ON : 27/Feb/2025 07:07AM
Lab No : 10132404	Sample Collected ON : 27/Feb/2025 07:08AM
Referred By : Dr.U1	Sample Received ON : 27/Feb/2025 07:59AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 27/Feb/2025 11:04AM
Doctor Advice : PP,FASTING,25 OH vit. D,HBA1C (EDTA),RF FACTOR	



Test Name	Result	Unit	Bio. Ref. Range	Method
FASTING				
Blood Sugar Fasting	150.2	mg/dl	70 - 110	Hexokinase
PP				
Blood Sugar PP	165.0	mg/dl	up to - 170	Hexokinase

*** End Of Report ***



CHARAK



MC-2491

Print.Date/Time: 27-02-2025 12:51:57

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sham

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)