


Patient Name : Ms.JAHIDA	Visit No : CHA250035649
Age/Gender : 61 Y/F	Registration ON : 27/Feb/2025 04: 25PM
Lab No : 10132945	Sample Collected ON : 27/Feb/2025 04: 28PM
Referred By : Dr.ESIC HOSPITAL LUCKNOW	Sample Received ON : 27/Feb/2025 04: 45PM
Refer Lab/Hosp : ESIC HOSPITAL LUCKNOW	Report Generated ON : 27/Feb/2025 08: 24PM
Doctor Advice : RF FACTOR,ANTI CCP TITRE,CRP (Quantitative)	



Test Name	Result	Unit	Bio. Ref. Range	Method
RF FACTOR				
RHEUMATOID FACTOR	20.10	IU/ml	0 - 14	

SUMMARY : Rheumatoid factors (RF) group of autoantibodies belonging to all immunoglobulin classes directed against the FC fragment of altered or complexed Igg .Diagnostic test for RF determination identify mainly RF of the IgM class which are detectable in several rheumatic diseases, mainly of inflammatory origin.

RF occur in approx 70 -80 % of patients with rheumatoid arthritis (RA), but they are not specific for RA as elevated concentrations are also observed in various non rheumatic disease & in approx 10 % of the elderly population without clinical symptoms of RA. High RF concentrations in RA are often associated with a more progressive clinical course of the disease .However,a positive RF value has to be confirmed by clinical & other laboratory findings.



CHARAK

[Checked By]

Print.Date/Time: 27-02-2025 21:04:02

*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADABKHAN
PATHOLOGIST

Dr. SYED SAIF AHMAD
MD (MICROBIOLOGY)

Patient Name : Ms.JAHIDA	Visit No : CHA250035649
Age/Gender : 61 Y/F	Registration ON : 27/Feb/2025 04: 25PM
Lab No : 10132945	Sample Collected ON : 27/Feb/2025 04: 28PM
Referred By : Dr.ESIC HOSPITAL LUCKNOW	Sample Received ON : 27/Feb/2025 04: 45PM
Refer Lab/Hosp : ESIC HOSPITAL LUCKNOW	Report Generated ON : 27/Feb/2025 08: 24PM
Doctor Advice : RF FACTOR,ANTI CCP TITRE,CRP (Quantitative)	



Test Name	Result	Unit	Bio. Ref. Range	Method
CRP-QUANTITATIVE				
CRP-QUANTITATIVE TEST	4.2	MG/L	0.1 - 6	

Method: Immunoturbidimetric

(Method: Immunoturbidimetric on photometry system)

SUMMARY : C - reactive protien (CRP) is the best known among the acute phase protiens, a group of protien whose concentration increases in blood as a response to inflammatory disorders.CRP is normally present in low concentration in blood of healthy individuals (< 1mg/L). It is elevated up to 500 mg/L in acute inflammatory processes associated with bacterial infections, post operative conditions tissue damage already after 6 hours reaching a peak at 48 hours.. The measurment of CRP represents a useful aboratory test for detection of acute infection as well as for monitoring inflammtory proceses also in acute rheumatic & gastrointestinal disease. In recent studies it has been shows that in apparently healthy subjects there is a direct orrelation between CRP concentrations & the risk of developing oronary heart disease (CHD).

hsCRP cut off for risk assessment as per CDC/AHA

Level	Risk
<1.0	Low
1.0-3.0	Average
>3.0	High

All reports to be clinically corelated

*** End Of Report ***

[Checked By]

Print.Date/Time: 27-02-2025 21:04:04

*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADABKHAN
PATHOLOGIST

Dr. SYED SAIF AHMAD
MD (MICROBIOLOGY)