

292/05, Tulsidas Marg, Basement Chowk, Lucknow-226 003

Phone: 0522-4062223, 9305548277, 8400888844 9415577933, 9336154100, Tollfree No.: 8688360360

E-mail: charak1984@gmail.com

CMO Reg. No. RMEE 2445133 NABLReg. No.MC-2491 Certificate No. MIS-2023-0218

Patient Name : Mr.NAVEEN KUMAR RATHOUR

Age/Gender : 42 Y/M Lab No : 10131914

Referred By : Dr.KGMU Refer Lab/Hosp : CHARAK NA

Doctor Advice : TB GOLD (3 metropolis tubes)

Visit No : CHA250034618

Registration ON : 26/Feb/2025 10:57AM

Sample Collected ON : 26/Feb/2025 10:58AM Sample Received ON : 27/Feb/2025 04:42PM

Report Generated ON : 28/Feb/2025 09:10AM

Test Name	Result	Unit	Bio. Ref. Range	Method	
TB GOLD					
Final Report	POSITIVE		NEGATIVE		
INTERFERON GAMMA IN NIL TUBE	0.50				
INTERFERON GAMMA IN ANTIGEN	2.62				
TUBE					
INTERFERON GAMMA TB (ANTIGEN	2.12				
MINUS NIL)					
INTERPRETATION					

NIL TUBE	INTERFE	RON GAMM	IA TB FINAL]	RESULT INTERPRETATION		
(IU/ML)	ANTIGEN MINUS NIL(IU/ML)					
	< 0.35		Negative	M.tuberculosis infection unlikely		
<=8.00	≥0.35 & <25%	% of Nil tube	Negative	M.tuberculosis infection unlikely		
	≥0.35 & ≥25%	6 of Nil tube	Positive	M.tuberculosis infection likely		
·			ĺ	This may be due to excessive level of		
>8.00	Any result		Indeterminate	circulating gamma interferon or		
		V.		presence of heterophile antibodies		

COMMENT:

A Quantiferon test is a test for latent Tuberculosis infection and helps in diagnosing M.tuberculosis complex infection including MOTT like Mkanasasii, M.szulgai and M.marinum.

An indeterminate result may occur with insufficient lymphocytes or inability of the patients lymphocytes to generate IFN-y response. This assay cannot differentiate between Latent infection and Active TB.

Magnitude of measured Gamma Interferon cannot be correlated with stage or degree of infection, level of immune responsiveness or likelihood ofprogression to active disease.

FALSE NEGATIVE RESULTS CAN BE THERE:

- I .False negative results maybe obtained if sample is taken prior to development of immune response. CDC recommends repeat test afir 8 - 10 weeks in case of high suspicion of tuberculosis.
- 2. Immunocompromised patients can also show false negative results.

*** End Of Report ***

