

Patient Name : Mr. IRSHAD KHAN	Visit No : CHA250036076
Age/Gender : 41 Y/M	Registration ON : 28/Feb/2025 11:18AM
Lab No : 10133372	Sample Collected ON : 28/Feb/2025 11:19AM
Referred By : Dr. SR SIDDIQUI	Sample Received ON : 28/Feb/2025 11:19AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 28/Feb/2025 02:07PM
Doctor Advice : URINE COM. EXMAMINATION, PSA-TOTAL	



Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

URINE EXAMINATION REPORT

Colour-U	YELLOW		Light Yellow	
Appearance (Urine)	CLEAR		Clear	
Specific Gravity	1.020		1.005 - 1.025	
pH-Urine	Acidic (6.0)		4.5 - 8.0	
PROTEIN	Absent	mg/dl	ABSENT	Dipstick
Glucose	Absent			
Ketones	Absent		Absent	
Bilirubin-U	Absent		Absent	
Blood-U	Absent		Absent	
Urobilinogen-U	0.20	EU/dL	0.2 - 1.0	
Leukocytes-U	Absent		Absent	
NITRITE	Absent		Absent	
MICROSCOPIC EXAMINATION				
Pus cells / hpf	5-6	/hpf	< 5/hpf	
Epithelial Cells	3-4	/hpf	0 - 5	
RBC / hpf	Nil		< 3/hpf	

CHARAK

[Checked By]



Sharma

DR. NISHANT SHARMA PATHOLOGIST DR. SHADAB PATHOLOGIST Dr. SYED SAIF AHMAD MD (MICROBIOLOGY)

Print.Date/Time: 28-02-2025 17:01:56

*Patient Identity Has Not Been Verified. Not For Medicolegal

Patient Name : Mr. IRSHAD KHAN	Visit No : CHA250036076
Age/Gender : 41 Y/M	Registration ON : 28/Feb/2025 11:18AM
Lab No : 10133372	Sample Collected ON : 28/Feb/2025 11:19AM
Referred By : Dr. SR SIDDIQUI	Sample Received ON : 28/Feb/2025 11:38AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 28/Feb/2025 04:24PM
Doctor Advice : URINE COM. EXMAMINATION, PSA-TOTAL	



Test Name	Result	Unit	Bio. Ref. Range	Method
PSA-TOTAL				
PROSTATE SPECIFIC ANTIGEN	0.83	ng/mL	0.2-4.0	CLIA

COMMENT : 1. Prostate specific antigen (PSA) is useful for diagnosis of disseminated CA prostate & its equential measurement is the most sensitive measure of monitoring treatment of disseminated CA prostate with its shorter half life (half life of 2.2 days only) it is superior to prostatic acis phosphatase(PAP). PSA is elevated in nearly all patients with stage D carcinoma whereas PAP is elevated in only 45 % of patient. Mild PSA elevation are also reported in some patients of BHP.

2. Blood samples should be obtained before prostate biopsy or prostatecomy or prostatic massage or digital pre rectal examination as it may result intrasient levation of PSA value for few days.

NOTE :- PSA values obtained in different types of PSA assay methods cannot be used interchangeably as the PSA value in a given sample varies with assays from different manufactures due to difference in assay methodology and reagent specificity. If in the course of monitoring a patient the assay method used for determination is changed, additional sequential testing should be carried out to confirm baseline value.

DONE BY;
Enhanced Chemiluminescence "VITROS ECI"

*** End Of Report ***

CHARAK



[Checked By]



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

Dr. Aditi D Agarwal
DR. ADITI D AGARWAL
PATHOLOGIST