

Patient Name : Mr.ASHOK SHUKLA
Age/Gender : 68 Y/M
Lab No : 10133810
Referred By : Dr.MANMEET SINGH
Refer Lab/Hosp : CHARAK NA
Doctor Advice : PSA-TOTAL
Visit No : CHA250036514
Registration ON : 28/Feb/2025 08:16PM
Sample Collected ON : 28/Feb/2025 08:18PM
Sample Received ON : 28/Feb/2025 09:52PM
Report Generated ON : 01/Mar/2025 09:30AM



Test Name	Result	Unit	Bio. Ref. Range	Method
PSA-TOTAL				
PROSTATE SPECIFIC ANTIGEN	3.09	ng/mL	0.2-4.0	CLIA

COMMENT : 1. Prostate specific antigen (PSA) is useful for diagnosis of disseminated CA prostate & its equential measurement is the most sensitive measure of monitoring treatment of disseminated CA prostate with its shorter half life (half life of 2.2 days only) it is superior to prostatic acis phosphatase(PAP). PSA is elevated in nearly all patients with stage D carcinoma whereas PAP is elevated in only 45 % of patient. Mild PSA elevation are also reported in some patients of BHP.
2. Blood samples should be obtained before prostate biopsy or prostatecomy or prostatic massage or digital pre rectal examination as it may result intrasient levation of PSA value for few days.

NOTE :- PSA values obtained in different types of PSA assay methods cannot be used interchangeably as the PSA value in a given sample varies with assays from different manufactures due to difference in assay methodology and reagent specificity. If in the course of monitoring a patient the assay method used for determination is changed, additional sequential testing should be carried out to confirm baseline value.

DONE BY;
Enhanced Chemiluminescence "VITROS ECI"

*** End Of Report ***

CHARAK



MC-2491

Print.Date/Time: 01-03-2025 10:05:11

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

Dr. SYED SAIF AHMAD
MD (MICROBIOLOGY)