

Patient Name : Ms.NISHA	Visit No : CHA250036923
Age/Gender : 22 Y/F	Registration ON : 01/Mar/2025 11:50AM
Lab No : 10134219	Sample Collected ON : 01/Mar/2025 11:52AM
Referred By : Dr.KGMU	Sample Received ON : 01/Mar/2025 11:58AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 01/Mar/2025 02:12PM
Doctor Advice : HBSAg,LFT,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
HEPATITIS B SURFACE ANTIGEN (HBsAg)				
Sample Type : SERUM				

HEPATITIS B SURFACE ANTIGEN	REACTIVE (4240)		<1 - Non Reactive >1 - Reactive	CMIA
-----------------------------	--------------------	--	------------------------------------	------

Note: This is only a Screening test. Confirmation of the result (Non Reactive/Reactive)should be done by performing a PCR based test.

COMMENTS:

- HBsAg is the first serological marker after infection with Hepatitis B Virus appearing one to ten weeks after exposure and two to eight weeks before the onset of clinical symptoms. HBsAg persists during the acute phase and clears late in the convalescence phase. Failure to clear HBsAg within six months indicates a chronic HBsAg carrier state. HBsAg assays are used to identify the persons infected with HBV and to prevent transmission of the virus by blood and blood products as well as to monitor the status of infected individuals in combination with other hepatitis B serological markers.
- Borderline cases must be confirmed with confirmatory neutralizing assay.

LIMITATIONS:

- Results should be used in conjunction with patient history and other hepatitis markers for diagnosis of acute and chronic infections.
- Specimens from patients who have received preparations of mouse monoclonal antibodies for diagnosis or therapy may contain human anti-mouse antibodies (HAMA) which may produce anomalous values when tested with assay kits that employs mouse monoclonal antibodies.
- Heterophilic antibodies in human serum can react with reagent immunoglobulins, interfering with in vitro immunoassays. Patients routinely exposed to animals or animal serum products can be prone to this interference and anomalous results may be observed.
- Cross reactivity for specimens from individual with medical conditions (Pregnancy, HIV etc) has been observed.
- HBsAg mutations may result in a false negative result in some HBsAg assays.
- If HBsAg results are inconsistent with clinical evidence, additional testing is suggested to confirm the result.

[Checked By]

Print.Date/Time: 01-03-2025 14:47:09

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Ms.NISHA	Visit No : CHA250036923
Age/Gender : 22 Y/F	Registration ON : 01/Mar/2025 11:50AM
Lab No : 10134219	Sample Collected ON : 01/Mar/2025 11:52AM
Referred By : Dr.KGMU	Sample Received ON : 01/Mar/2025 11:58AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 01/Mar/2025 02:12PM
Doctor Advice : HBSAg,LFT,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
CBC (COMPLETE BLOOD COUNT)				
Hb	11.3	g/dl	12 - 15	Non Cyanide
R.B.C. COUNT	4.00	mil/cmm	3.8 - 4.8	Electrical Impedence
PCV	35.6	%	36 - 45	Pulse hieght detection
MCV	88.1	fL	80 - 96	calculated
MCH	28.0	pg	27 - 33	Calculated
MCHC	31.7	g/dL	30 - 36	Calculated
RDW	15.1	%	11 - 15	RBC histogram derivation
RETIC	0.9 %	%	0.5 - 2.5	Microscopy
TOTAL LEUCOCYTES COUNT	5790	/cmm	4000 - 10000	Flocytrometry
DIFFERENTIAL LEUCOCYTE COUNT				
NEUTROPHIL	54	%	40 - 75	Flowcytometry
LYMPHOCYTES	40	%	25 - 45	Flowcytometry
EOSINOPHIL	3	%	1 - 6	Flowcytometry
MONOCYTE	3	%	2 - 10	Flowcytometry
BASOPHIL	0	%	00 - 01	Flowcytometry
PLATELET COUNT	163,000	/cmm	150000 - 450000	Elect Imped..
PLATELET COUNT (MANUAL)	163000	/cmm	150000 - 450000	Microscopy .
Absolute Neutrophils Count	3,127	/cmm	2000 - 7000	Calculated
Absolute Lymphocytes Count	2,316	/cmm	1000-3000	Calculated
Absolute Eosinophils Count	174	/cmm	20-500	Calculated
Absolute Monocytes Count	174	/cmm	200-1000	Calculated
Mentzer Index	22			
Peripheral Blood Picture	:			

Red blood cells are normocytic normochromic. Platelets are adequate. No immature cells or parasite seen.



[Checked By]



Sham


DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Ms.NISHA	Visit No : CHA250036923
Age/Gender : 22 Y/F	Registration ON : 01/Mar/2025 11:50AM
Lab No : 10134219	Sample Collected ON : 01/Mar/2025 11:52AM
Referred By : Dr.KGMU	Sample Received ON : 01/Mar/2025 11:58AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 01/Mar/2025 12:48PM
Doctor Advice : HBSAg,LFT,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
LIVER FUNCTION TEST				
TOTAL BILIRUBIN	1.12	mg/dl	0.4 - 1.1	Diazonium Ion
CONJUGATED (D. Bilirubin)	0.23	mg/dL	0.00-0.30	Diazotization
UNCONJUGATED (I.D. Bilirubin)	0.89	mg/dL	0.1 - 1.0	Calculated
ALK PHOS	164.20	U/L	30 - 120	PNPP, AMP Buffer
SGPT	22.0	U/L	5 - 40	UV without P5P
SGOT	24.0	U/L	5 - 40	UV without P5P

*** End Of Report ***



CHARAK



[Checked By]



Sharma

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)