


Patient Name : Mr.SHVI KUMAR GUPTA
Age/Gender : 60 Y/M
Lab No : 10134735
Referred By : SELF
Refer Lab/Hosp : CHARAK NA
Doctor Advice : PP,FASTING,T3T4TSH

Visit No : CHA250037440
Registration ON : 02/Mar/2025 08:16AM
Sample Collected ON : 02/Mar/2025 08:17AM
Sample Received ON : 02/Mar/2025 08:36AM
Report Generated ON : 02/Mar/2025 02:19PM



Test Name	Result	Unit	Bio. Ref. Range	Method
FASTING				
Blood Sugar Fasting	97.9	mg/dl	70 - 110	Hexokinase
PP				
Blood Sugar PP	124.0	mg/dl	up to - 170	Hexokinase



CHARAK



[Checked By]



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

Dr. Aditi D. Agarwal
DR. ADITI D AGARWAL
PATHOLOGIST

Patient Name : Mr.SHVI KUMAR GUPTA	Visit No : CHA250037440
Age/Gender : 60 Y/M	Registration ON : 02/Mar/2025 08:16AM
Lab No : 10134735	Sample Collected ON : 02/Mar/2025 08:17AM
Referred By : SELF	Sample Received ON : 02/Mar/2025 08:36AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 02/Mar/2025 10:06AM
Doctor Advice : PP,FASTING,T3T4TSH	



Test Name	Result	Unit	Bio. Ref. Range	Method
T3T4TSH				
T3	2.36	nmol/L	1.49-2.96	ECLIA
T4	160.39	n mol/l	63 - 177	ECLIA
TSH	4.08	uIU/ml	0.47 - 4.52	ECLIA

Note

- (1) Patients having low T3 & T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or autoimmune disorders.
- (2) Patients having low T3 & T4 levels but high TSH levels suffer from grave's disease, toxic adenoma or sub-acute thyroiditis.
- (3) Patients having either low or normal T3 & T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.
- (4) Patients having high T3 & T4 levels but normal TSH levels may suffer from toxic multinodular goitre. This condition is mostly asymptomatic and may cause transient hyperthyroidism but no persistent symptoms.
- (5) Patient with high or normal T3 & T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 Toxicosis respectively.
- (6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.
- (7) There are many drugs for eg. Glucocorticoids, dopamine, Lithium, iodides, oral radiographic dyes, etc. Which may affect the thyroid function tests.
- (8) Generally when total T3 & T4 results are indecisive then Free T3 & Free T4 test are recommended for further confirmation along with

(1 Beckman Dxi-600 2. ELECTRO-CHEMILUMINESCENCE TECHNIQUE BY ELECSYS -E411)

*** End Of Report ***

CHARAK



[Checked By]



Sham

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)