

Patient Name	: Mr.ADNAN YUSUF	Visit No	: CHA250037672
Age/Gender	: 50 Y/M	Registration ON	: 02/Mar/2025 12:23PM
Lab No	: 10134967	Sample Collected ON	: 02/Mar/2025 12:23PM
Referred By	: Dr.AK TRIPATHI	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 02/Mar/2025 03:24PM

COLOUR DOPPLER STUDY OF BILATERAL LOWER LIMB VEINS

- Bilateral common femoral, superficial femoral, popliteal and visualized parts of bilateral tibial veins reveal clear lumen and normal colour flow with normal phasicity, compressibility and augmentation response.
- *Bilateral anterior and posterior tibial veins could not be very well evaluated in complete extent.*
- There is maintained color flow across bilateral sapheno-femoral junctions, competence could be assessed as patient was unable to perform valsalva maneuver.
- *Bilateral sapheno popliteal junctions could not be very well evaluated.*

IMPRESSION:

- **NO EVIDENCE OF DEEP VEIN THROMBOSIS IN VISUALIZED VEINS.**

CHARAK

Clinical correlation is necessary.

[DR. JAYENDRA K. ARYA, MD]

*** End Of Report ***

