

Patient Name	: Ms.SAVITRI	Visit No	: CHA250037909
Age/Gender	: 27 Y/F	Registration ON	: 03/Mar/2025 09:10AM
Lab No	: 10135204	Sample Collected ON	: 03/Mar/2025 09:10AM
Referred By	: Dr.ESIC HOSPITAL LUCKNOW	Sample Received ON	:
Refer Lab/Hosp	: ESIC HOSPITAL LUCKNOW	Report Generated ON	: 03/Mar/2025 11:08AM

NCCT STUDY OF PARANASAL SINUSES

- Nasal septum is deviated with convexity towards left side with the septal spur impending upon left inferior nasal turbinate.
- Mucosal thickening is seen in left maxillary ,ethmoid and frontal sinuses with occlusion of left osteomeatal complex and fronto-ethmoidal recess.
- Small right concha bullosa is seen .
- Right, right frontal and bilateral sphenoid sinuses are clear.
- Minimal mucosal thickening is seen in right ethmoid sinus .
- Minimal to mild mucosal thickening is seen involving bilateral [left > right] nasal turbinates .
- No mass lesion is seen in nasopharynx or nasal cavity.

IMPRESSION:

- FINDINGS ARE SUGGESTIVE OF LEFT MAXILLARY , ETHMOID AND FRONTAL SINUSITIS.
- MINIMAL MUCOSAL THICKENING IN RIGHT ETHMOID SINUS .
- DEVIATED NASAL SEPTUM TOWARDS LEFT SIDE.

Clinical correlation is necessary.

[DR. JAYENDRA KUMAR, MD]

TRANSCRIBED BY: ANUP

*** End Of Report ***

