

Patient Name : Mr. SUBHASH CHANDRA YADAV Visit No : CHA250038445
Age/Gender : 38 Y/M Registration ON : 03/Mar/2025 05:21PM
Lab No : 10135740 Sample Collected ON : 03/Mar/2025 05:21PM
Referred By : Dr. ESIC HOSPITAL LUCKNOW Sample Received ON :
Refer Lab/Hosp : ESIC HOSPITAL LUCKNOW Report Generated ON : 03/Mar/2025 07:20PM

MRI: PERINEUM

IMAGING SEQUENCES (NCMR)

AXIAL: T1, T2 & T2 FS Wis. **CORONAL:** T2 FS Wis. **SAGITTAL:** TIRM

There is a linear T2 hyperintense tract (approx 28mm in length) extending from the external opening in left perianal region (at 6 O' clock position just away from the anal verge) through the intersphincteric plane to the posterior wall of anal canal at 6 O'clock position (approx 21mm above the anal verge). Another small T2 hyperintense tract (approx 15mm in length) is seen extending from left posterior perianal region at 4 O'clock position through the intersphincteric plane and indenting the left lateral wall of anal canal at 3 O'clock position, just above the anal verge. Bilateral ischiorectal fossae are normal.

Urinary bladder is well distended.

Prostate is normal in outline and signal intensity.

Both the seminal vesicles are normal in size, outline and morphology.

No significant pelvic lymphadenopathy is noted.

Common iliac, external iliac and internal iliac vessels are showing normal flow void.

IMPRESSION

- **Features are suggestive of grade-I intersphincteric tracts.**

Please correlate clinically.

DR. RAVENDRA SINGH
MD

Typed by Ranjeet

*** End Of Report ***

