

Patient Name : Mr.RAM SEWAK	Visit No : CHA250039646
Age/Gender : 72 Y 1 D/M	Registration ON : 05/Mar/2025 11:48AM
<b>Lab No : 10136941</b>	Sample Collected ON : 05/Mar/2025 11:50AM
Referred By : Dr.RDSO LUCKNOW	Sample Received ON : 05/Mar/2025 12:03PM
Refer Lab/Hosp : RDSO LUCKNOW	Report Generated ON : 05/Mar/2025 01:25PM
Doctor Advice : PRO-BNP,PSA-TOTAL,NA+K+,CHEST PA	




Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

**PRO-BNP**

BNP (B type Natriuretic Peptide) 528.00

EXPECTED VALUES :- Expected values for blood donors (97.5 percentile)

	< 50years	N	> 50 years	N
WOMEN :	155 pg/ml	887	222pg/ml	248
MEN :	84 pg/ml	1381	194 pg/ml	432



**CHARAK**

[Checked By]

Print.Date/Time: 05-03-2025 14:15:18

\*Patient Identity Has Not Been Verified. Not For Medicolegal



*Sharma*

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD  
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Mr.RAM SEWAK	Visit No : CHA250039646
Age/Gender : 72 Y 1 D/M	Registration ON : 05/Mar/2025 11:48AM
<b>Lab No : 10136941</b>	Sample Collected ON : 05/Mar/2025 11:50AM
Referred By : Dr.RDSO LUCKNOW	Sample Received ON : 05/Mar/2025 12:03PM
Refer Lab/Hosp : RDSO LUCKNOW	Report Generated ON : 05/Mar/2025 01:25PM
Doctor Advice : PRO-BNP,PSA-TOTAL,NA+K+,CHEST PA	



Test Name	Result	Unit	Bio. Ref. Range	Method
<b>NA+K+</b>				
SODIUM Serum	<b>133.0</b>	MEq/L	135 - 155	ISE Direct
POTASSIUM Serum	5.2	MEq/L	3.5 - 5.5	ISE Direct
<b>PSA-TOTAL</b>				
PROSTATE SPECIFIC ANTIGEN	<b>5.2</b>	ng/mL	0.2-4.0	CLIA

COMMENT : 1. Prostate specific antigen (PSA) is useful for diagnosis of disseminated CA prostate & its sequential measurement is the most sensitive measure of monitoring treatment of disseminated CA prostate with its shorter half life (half life of 2.2 days only) it is superior to prostatic acid phosphatase(PAP). PSA is elevated in nearly all patients with stage D carcinoma whereas PAP is elevated in only 45 % of patient. Mild PSA elevation are also reported in some patients of BHP.

2. Blood samples should be obtained before prostate biopsy or prostatectomy or prostatic massage or digital pre rectal examination as it may result in transient elevation of PSA value for few days.

NOTE :- PSA values obtained in different types of PSA assay methods cannot be used interchangeably as the PSA value in a given sample varies with assays from different manufactures due to difference in assay methodology and reagent specificity. If in the course of monitoring a patient the assay method used for determination is changed, additional sequential testing should be carried out to confirm baseline value.

DONE BY;  
Enhanced Chemiluminescence "VITROS ECI"

\*\*\* End Of Report \*\*\*



[Checked By]



*Sharma*

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD  
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Mr.RAM SEWAK Visit No : CHA250039646  
Age/Gender : 72 Y 1 D/M Registration ON : 05/Mar/2025 11:48AM  
**Lab No : 10136941** Sample Collected ON : 05/Mar/2025 11:48AM  
Referred By : Dr.RDSO LUCKNOW Sample Received ON :  
Refer Lab/Hosp : RDSO LUCKNOW Report Generated ON : 05/Mar/2025 01:41PM

**SKIAGRAM CHEST PA VIEW**

- Fibrotic opacities are seen in both mid zones.
- Both lung fields show hyperinflation and increased lucency in lower zone.
- Bilateral hilar shadows are prominent.
- Cardiac shadow is within normal limits.
- Both CP angles are blunted.
- Soft tissue and bony cage are seen normally.
- Both domes of diaphragm are slightly depressed.

**IMPRESSION:**

- **FINDINGS ARE SUGGESTIVE EARLY C.O.P.D. CHANGES.**

*To be correlated with previous records.*

**Clinical correlation is necessary.**

**[DR. RAJESH KUMAR SHARMA, MD]**

TRANSCRIBED BY: ANUP

\*\*\* End Of Report \*\*\*

