

Patient Name : Mr.SAFDAR HUSAIN	Visit No : CHA250040184
Age/Gender : 72 Y 4 M 13 D/M	Registration ON : 06/Mar/2025 06: 54AM
Lab No : 10137479	Sample Collected ON : 06/Mar/2025 06: 57AM
Referred By : Dr.KAUSER USMAN	Sample Received ON : 06/Mar/2025 07: 11AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 06/Mar/2025 09: 18AM
Doctor Advice : T3T4TSH,FASTING	



Test Name	Result	Unit	Bio. Ref. Range	Method
FASTING				
Blood Sugar Fasting	127.8	mg/dl	70 - 110	Hexokinase



[Checked By]



Sharma

DR. NISHANT SHARMA PATHOLOGIST DR. SHADAB PATHOLOGIST Dr. SYED SAIF AHMAD MD (MICROBIOLOGY)

Patient Name : Mr.SAFDAR HUSAIN	Visit No : CHA250040184
Age/Gender : 72 Y 4 M 13 D/M	Registration ON : 06/Mar/2025 06: 54AM
Lab No : 10137479	Sample Collected ON : 06/Mar/2025 06: 57AM
Referred By : Dr.KAUSER USMAN	Sample Received ON : 06/Mar/2025 07: 11AM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 06/Mar/2025 09: 18AM
Doctor Advice : T3T4TSH,FASTING	



Test Name	Result	Unit	Bio. Ref. Range	Method
T3T4TSH				
T3	2.31	nmol/L	1.49-2.96	ECLIA
T4	170.77	n mol/l	63 - 177	ECLIA
TSH	2.25	uIU/ml	0.47 - 4.52	ECLIA

Note

- (1) Patients having low T3 & T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or autoimmune disorders.
- (2) Patients having low T3 & T4 levels but high TSH levels suffer from grave's disease, toxic adenoma or sub-acute thyroiditis.
- (3) Patients having either low or normal T3 & T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.
- (4) Patients having high T3 & T4 levels but normal TSH levels may suffer from toxic multinodular goitre. This condition is mostly asymptomatic and may cause transient hyperthyroidism but no persistent symptoms.
- (5) Patient with high or normal T3 & T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 Toxicosis respectively.
- (6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.
- (7) There are many drugs for eg. Glucocorticoids, dopamine, Lithium, iodides, oral radiographic dyes, etc. Which may affect the thyroid function tests.
- (8) Generally when total T3 & T4 results are indecisive then Free T3 & Free T4 test are recommended for further confirmation along with

(1 Beckman Dxi-600 2. ELECTRO-CHEMILUMINESCENCE TECHNIQUE BY ELECSYS -E411)

*** End Of Report ***

CHARAK



[Checked By]



Sharma

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)