

Patient Name : Ms.RUBINA KHATOON	Visit No : CHA250040845
Age/Gender : 33 Y/F	Registration ON : 06/Mar/2025 07:55PM
<b>Lab No : 10138140</b>	Sample Collected ON : 06/Mar/2025 07:58PM
Referred By : Dr.ANSHUMALA RASTOGI	Sample Received ON : 06/Mar/2025 07:58PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 07/Mar/2025 10:17AM
Doctor Advice : URINE COM. EXMAMINATION,TSH,HCV,HBSAg,HIV,VDRL,RANDOM,BLOOD GROUP,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
<b>BLOOD GROUP</b>				
Blood Group	"B"			
Rh (Anti -D)	Negative			

<b>HEPATITIS B SURFACE ANTIGEN (HBsAg)</b>				
Sample Type : SERUM				
HEPATITIS B SURFACE ANTIGEN	NON REACTIVE		<1 - Non Reactive >1 - Reactive	CMIA

Note: This is only a Screening test. Confirmation of the result ( Non Reactive/Reactive)should be done by performing a PCR based test.

**COMMENTS:**

- HBsAg is the first serological marker after infection with Hepatitis B Virus appearing one to ten weeks after exposure and two to eight weeks before the onset of clinical symptoms. HBsAg persists during the acute phase and clears late in the convalescence phase. Failure to clear HBsAg within six months indicates a chronic HBsAg carrier state. HBsAg assays are used to identify the persons infected with HBV and to prevent transmission of the virus by blood and blood products as well as to monitor the status of infected individuals in combination with other hepatitis B serological markers.
- Borderline cases must be confirmed with confirmatory neutralizing assay.

**LIMITATIONS:**

- Results should be used in conjunction with patient history and other hepatitis markers for diagnosis of acute and chronic infections.
- Specimens from patients who have received preparations of mouse monoclonal antibodies for diagnosis or therapy may contain human anti-mouse antibodies (HAMA) which may produce anomalous values when tested with assay kits that employs mouse monoclonal antibodies.
- Heterophilic antibodies in human serum can react with reagent immunoglobulins, interfering with in vitro immunoassays. Patients routinely exposed to animals or animal serum products can be prone to this interference and anomalous results may be observed.
- Cross reactivity for specimens from individual with medical conditions (Pregnancy, HIV etc) has been observed.
- HBsAg mutations may result in a false negative result in some HBsAg assays.
- If HBsAg results are inconsistent with clinical evidence, additional testing is suggested to confirm the result.

[Checked By]

Print.Date/Time: 07-03-2025 10:50:12

\*Patient Identity Has Not Been Verified. Not For Medicolegal



*Sharma*

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD  
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Ms.RUBINA KHATOON	Visit No : CHA250040845
Age/Gender : 33 Y/F	Registration ON : 06/Mar/2025 07: 55PM
<b>Lab No : 10138140</b>	Sample Collected ON : 06/Mar/2025 07: 58PM
Referred By : Dr.ANSHUMALA RASTOGI	Sample Received ON : 06/Mar/2025 07: 58PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 07/Mar/2025 10: 17AM
Doctor Advice : URINE COM. EXMAMINATION,TSH,HCV,HBSAg,HIV,VDRL,RANDOM,BLOOD GROUP,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

**HIV**

HIV-SEROLOGY	NON REACTIVE	<1.0 : NON REACTIVE >1.0 : REACTIVE
--------------	--------------	--

Done by: Vitros ECI ( Sandwich Assay)

Note:-Elisa test is a screening method for HIV.It is known to give false Positive & Negative result.  
Hence confirmation:"Western Blot" method is advised.

**HCV**

Anti-Hepatitis C Virus Antibodies.	NON REACTIVE	< 1.0 : NON REACTIVE > 1.0 : REACTIVE	Sandwich Assay
------------------------------------	--------------	--	----------------

Done by: Vitros ECI ( Sandwich Assay)

Note:This is only a Screening test. Confirmation of the result ( Non Reactive/Reactive)should be done by performing a PCR based test.

**VDRL**

VDRL	NON REACTIVE	Slide Agglutination
------	--------------	---------------------

**URINE EXAMINATION REPORT**

Colour-U	YELLOW	Light Yellow	
Appearance (Urine)	CLEAR	Clear	
Specific Gravity	<b>1.015</b>	1.005 - 1.025	
pH-Urine	Acidic (6.0)	4.5 - 8.0	
PROTEIN	Absent	ABSENT	Dipstick
Glucose	Absent		
Ketones	Absent	Absent	
Bilirubin-U	Absent	Absent	
Blood-U	Absent	Absent	
Urobilinogen-U	0.20	EU/dL	0.2 - 1.0
Leukocytes-U	Absent	Absent	
NITRITE	Absent	Absent	

**MICROSCOPIC EXAMINATION**

Pus cells / hpf	Nil	/hpf	< 5/hpf
Epithelial Cells	3-4	/hpf	0 - 5
RBC / hpf	Nil		< 3/hpf

[Checked By]

Print.Date/Time: 07-03-2025 10:50:13

\*Patient Identity Has Not Been Verified. Not For Medicolegal



*Sharma*

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD  
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Ms.RUBINA KHATOON	Visit No : CHA250040845
Age/Gender : 33 Y/F	Registration ON : 06/Mar/2025 07:55PM
<b>Lab No : 10138140</b>	Sample Collected ON : 06/Mar/2025 07:58PM
Referred By : Dr.ANSHUMALA RASTOGI	Sample Received ON : 06/Mar/2025 10:26PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 07/Mar/2025 09:18AM
Doctor Advice : URINE COM. EXMAMINATION,TSH,HCV,HBSAg,HIV,VDRL,RANDOM,BLOOD GROUP,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
<b>CBC (COMPLETE BLOOD COUNT)</b>				
Hb	13.0	g/dl	12 - 15	Non Cyanide
R.B.C. COUNT	4.60	mil/cmm	3.8 - 4.8	Electrical Impedence
PCV	39.2	%	36 - 45	Pulse hieght detection
MCV	85.2	fL	80 - 96	calculated
MCH	28.3	pg	27 - 33	Calculated
MCHC	33.2	g/dL	30 - 36	Calculated
RDW	13.1	%	11 - 15	RBC histogram derivation
RETIC	0.7 %	%	0.5 - 2.5	Microscopy
TOTAL LEUCOCYTES COUNT	9990	/cmm	4000 - 10000	Flocytrometry
<b>DIFFERENTIAL LEUCOCYTE COUNT</b>				
NEUTROPHIL	73	%	40 - 75	Flowcytometry
LYMPHOCYTES	<b>23</b>	%	25 - 45	Flowcytometry
EOSINOPHIL	<b>0</b>	%	1 - 6	Flowcytometry
MONOCYTE	4	%	2 - 10	Flowcytometry
BASOPHIL	<b>0</b>	%	00 - 01	Flowcytometry
PLATELET COUNT	191,000	/cmm	150000 - 450000	Elect Imped..
PLATELET COUNT (MANUAL)	191000	/cmm	150000 - 450000	Microscopy .
Absolute Neutrophils Count	<b>7,293</b>	/cmm	2000 - 7000	Calculated
Absolute Lymphocytes Count	2,298	/cmm	1000-3000	Calculated
Absolute Monocytes Count	400	/cmm	200-1000	Calculated
Mentzer Index	19			
Peripheral Blood Picture	:			

Red blood cells are normocytic normochromic. Platelets are adequate. No immature cells or parasite seen.



[Checked By]



*Sham*

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD  
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Ms.RUBINA KHATOON	Visit No : CHA250040845
Age/Gender : 33 Y/F	Registration ON : 06/Mar/2025 07:55PM
<b>Lab No : 10138140</b>	Sample Collected ON : 06/Mar/2025 07:58PM
Referred By : Dr.ANSHUMALA RASTOGI	Sample Received ON : 06/Mar/2025 08:07PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 07/Mar/2025 09:11AM
Doctor Advice : URINE COM. EXMAMINATION,TSH,HCV,HBSAg,HIV,VDRL,RANDOM,BLOOD GROUP,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
<b>BLOOD SUGAR RANDOM</b>				
BLOOD SUGAR RANDOM	125	mg/dl	70 - 170	Hexokinase

<b>TSH</b>				
TSH	1.90	uIU/ml	0.47 - 4.52	ECLIA

**Note**

- (1) Patients having low T3 & T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or autoimmune disorders.
- (2) Patients having low T3 & T4 levels but high TSH levels suffer from grave's disease, toxic adenoma or sub-acute thyroiditis.
- (3) Patients having either low or normal T3 & T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.
- (4) Patients having high T3 & T4 levels but normal TSH levels may suffer from toxic multinodular goitre. This condition is mostly asymptomatic and may cause transient hyperthyroidism but no persistent symptoms.
- (5) Patient with high or normal T3 & T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 Toxicosis respectively.
- (6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.
- (7) There are many drugs for eg. Glucocorticoids, dopamine, Lithium, iodides, oral radiographic dyes, etc. Which may affect the thyroid function tests.
- (8) Generally when total T3 & T4 results are indecisive then Free T3 & Free T4 test are recommended for further confirmation along with  
( 1 Beckman Dxi-600 2. ELECTRO-CHEMILUMINESCENCE TECHNIQUE BY ELECSYSYS -E411 )

\*\*\* End Of Report \*\*\*



[Checked By]



*Sham*

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD  
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)