

Patient Name	: Master ABBAS HAIDER	Visit No	: CHA250040951
Age/Gender	: 4 Y 5 M 0 D /M	Registration ON	: 07/Mar/2025 09:11AM
Lab No	: 10138246	Sample Collected ON	: 07/Mar/2025 09:11AM
Referred By	: Dr.NISHA PANDEY	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 07/Mar/2025 09:59AM

ULTRASOUND STUDY OF WHOLE ABDOMEN

- **Liver** is mildly enlarged in size (~125mm) and shows homogenous echotexture of liver parenchyma. No intrahepatic biliary radicle dilatation is seen. No space occupying lesion is seen. Hepatic veins and IVC are seen normally.
- **Gall bladder** is normal in size and shows anechoic lumen. No calculus / mass lesion is seen. GB walls are not thickened.
- **CBD** is normal at porta. No obstructive lesion is seen.
- **Portal vein** Portal vein is normal at porta.
- **Pancreas** is normal in size and shows homogenous echotexture of parenchyma. PD is not dilated. No parenchymal calcification is seen. No peripancreatic collection is seen.
- **Spleen** is normal in size and shows homogenous echotexture of parenchyma. No SOL is seen.
- **Few subcentimeteric mesenteric lymphnodes are seen measuring upto approx 11 x 7.1mm with maintained hilum.**
- No ascites is seen.
- **Both kidneys** are normal in size and position. No hydronephrosis is seen. No calculus or mass lesion is seen. Cortico-medullary differentiation is well maintained. Parenchymal thickness is normal. No scarring is seen. Right kidney measures 66 x 28 mm in size. Left kidney measures 68 x 26 mm in size.
- **Ureters** Both ureters are not dilated. UVJ are seen normally.
- **Urinary bladder** is normal in contour with anechoic lumen. No calculus or mass lesion is seen. UB walls are not thickened.
- Prostate is normal for child age group.
- **A target pattern bowel loop is seen in right para-umbilical region with no proximal bowel dilatation & the lesion is disappears after peristaltic movement-? Transient intussusception.**

OPINION:

- MILD HEPATOMEGALY.
- FEW SUBCENTIMETERIC MESENTERIC LYMPHNODES.
- A TARGET PATTERN BOWEL LOOP IN RIGHT PARA-UMBILICAL REGION WITH NO PROXIMAL BOWEL DILATATION & THE LESION IS DISAPPEARS AFTER PERISTALTIC MOVEMENT-? TRANSIENT INTUSSUSCEPTION.

Clinical correlation is necessary.

(DR. R.K. SINGH, MD)

Transcribed by Gausiya



Patient Name	: Master ABBAS HAIDER	Visit No	: CHA250040951
Age/Gender	: 4 Y 5 M 0 D /M	Registration ON	: 07/Mar/2025 09: 11AM
Lab No	: 10138246	Sample Collected ON	: 07/Mar/2025 09: 11AM
Referred By	: Dr.NISHA PANDEY	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 07/Mar/2025 09: 59AM

*** End Of Report ***

