

Patient Name : Mr.SIYA RAM VERMA  
Age/Gender : 66 Y/M  
**Lab No : 10138972**  
Referred By : Dr.KRISHNA KUMAR MITRA (CGHS)  
Refer Lab/Hosp : CGHS (BILLING)  
Doctor Advice : AFP,CEA,CA 19-9

Visit No : CHA250041677  
Registration ON : 08/Mar/2025 10:43AM  
Sample Collected ON : 08/Mar/2025 10:46AM  
Sample Received ON : 08/Mar/2025 11:04AM  
Report Generated ON : 08/Mar/2025 12:03PM



Test Name	Result	Unit	Bio. Ref. Range	Method
ALPHA-FETOPROTEIN (AFP)				
AFP	4.10	IU/ml	Upto 11.3	



**CHARAK**

[Checked By]

Print.Date/Time: 08-03-2025 16:50:08

\*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA  
PATHOLOGIST

DR. SHADAB  
PATHOLOGIST

Dr. SYED SAIF AHMAD  
MD (MICROBIOLOGY)

Patient Name : Mr.SIYA RAM VERMA	Visit No : CHA250041677
Age/Gender : 66 Y/M	Registration ON : 08/Mar/2025 10:43AM
<b>Lab No : 10138972</b>	Sample Collected ON : 08/Mar/2025 10:46AM
Referred By : Dr.KRISHNA KUMAR MITRA (CGHS)	Sample Received ON : 08/Mar/2025 11:04AM
Refer Lab/Hosp : CGHS (BILLING)	Report Generated ON : 08/Mar/2025 04:15PM
Doctor Advice : AFP,CEA,CA 19-9	



Test Name	Result	Unit	Bio. Ref. Range	Method
<b>CARCINOEMBRYONIC ANTIGEN (CEA)</b>				
CARCINOEMBRYONIC ANTIGEN (CEA)	1.80	ng/ml	0.00 - 4.50	

By.Electrochemiluminescence Immunoassay (ECLIA)

COMMENTS : CEA was first presented as a specific antigen for adenocarcinoma of the colon. More recent studies have demonstrated CEA presence in a variety of malignancies, particularly those involving ectodermal tissue of gastrointestinal or pulmonary origin. Small amounts have also been demonstrated in secretion of the colonic mucosa. Additionally, CEA like substances have been reported in normal bile from non-icteric patients.

CEA testing can have significant value in the monitoring of cancer patients. Persistent elevation in circulating CEA following treatment is strongly indicative of occult metastatic and / or residual disease. Also a persistent rising CEA value may be associated with progressive malignant disease or poor therapeutic response. A declining CEA value is generally indicative of favorable prognosis and good response to treatment. Measurement of CEA has been shown to be clinically relevant in the follow-up management of patients with colorectal, breast, lung, prostatic, pancreatic, ovarian, & a variety of other carcinomas suggest that the preoperative CEA level has prognostic significance. CEA testing is not recommended as a screening procedure to detect cancer in the general population, however, use of the CEA test as an adjunctive test in the prognosis & management of cancer patients has been widely accepted.

\*\*\* End Of Report \*\*\*

CHARAK

[Checked By]

Print.Date/Time: 08-03-2025 16:50:10

\*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA  
PATHOLOGIST

DR. SHADAB  
PATHOLOGIST

DR. ADITI D AGARWAL  
PATHOLOGIST

*Aditi D. Agarwal*