

Patient Name : Mr.MOHD MOID	Visit No : CHA250043102
Age/Gender : 50 Y/M	Registration ON : 10/Mar/2025 01:35PM
Lab No : 10140397	Sample Collected ON : 10/Mar/2025 01:36PM
Referred By : Dr.KGMU	Sample Received ON : 10/Mar/2025 01:43PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 10/Mar/2025 02:45PM
Doctor Advice : CREATININE,CT ANGIO ONE PART	



Test Name	Result	Unit	Bio. Ref. Range	Method
SERUM CREATININE				
CREATININE	0.70	mg/dl	0.50 - 1.40	Alkaline picrate-kinetic

*** End Of Report ***



CHARAK



[Checked By]



Sham

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name	: Mr.MOHD MOID	Visit No	: CHA250043102
Age/Gender	: 50 Y/M	Registration ON	: 10/Mar/2025 01:35PM
Lab No	: 10140397	Sample Collected ON	: 10/Mar/2025 01:35PM
Referred By	: Dr.KGMU	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 10/Mar/2025 07:01PM

CT ANGIOGRAPHY OF BILATERAL LOWER LIMBS

- Visualized abdominal aorta and bilateral iliac arteries show mild atherosclerotic changes with maintained post contrast opacification without obvious filling defect /significant luminal narrowing.
- Bilateral common femoral & superficial femoral arteries show mild atherosclerotic changes with maintained post contrast opacification without obvious filling defect /significant luminal narrowing.
- Bilateral profunda femoris arteries show maintained post contrast opacification without obvious filling defect /significant luminal narrowing
- Bilateral popliteal arteries appear normal in caliber with maintained post contrast opacification without obvious filling defect / significant luminal narrowing.
- Bilateral anterior and posterior tibial arteries appear normal in caliber with maintained post contrast opacification without obvious filling defect / significant luminal narrowing.
- Bilateral peroneal arteries show luminal attenuation in distal 1/3rd, however rest of the segments show maintained post contrast opacification without obvious filling defect.
- Bilateral dorsalis pedis arteries appear normal in caliber with maintained post contrast opacification without obvious filling defect / significant luminal narrowing.

OPINION:

- **MILD ATHEROSCLEROTIC CHANGES IN BILATERAL LOWER LIMB ARTERIES WITHOUT SIGNIFICANT LUMINAL NARROWING.**

Clinical correlation is necessary

[DR. JAYENDRA KUMAR, MD]

Transcribed by Rachna

*** End Of Report ***

