


Patient Name : Mr.TAJUDDIN	Visit No : CHA250043194
Age/Gender : 55 Y/M	Registration ON : 10/Mar/2025 02: 56PM
Lab No : 10140489	Sample Collected ON : 10/Mar/2025 02: 57PM
Referred By : Dr.YUSUF ANSARI	Sample Received ON : 10/Mar/2025 03: 21PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 10/Mar/2025 04: 52PM
Doctor Advice : CPK - MB,TROPONIN-T hs Stat,2D ECHO	



Test Name	Result	Unit	Bio. Ref. Range	Method
CPK-MB				
CPK-MB	2.37	U/L	Less than 25	

INTERPRETATION:

CK-MB is the enzyme being used as the definitive serum marker for the diagnosis of acute myocardial infarction. CK-MB, released after AMI, is detectable in blood as early as 3-4 hours after the onset of symptoms and remains elevated for approximately 65 hours post infarct.



CHARAK

[Checked By]

Print.Date/Time: 10-03-2025 18:17:47

*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

Aditi D Agarwal
DR. ADITI D AGARWAL
PATHOLOGIST

Patient Name : Mr.TAJUDDIN	Visit No : CHA250043194
Age/Gender : 55 Y/M	Registration ON : 10/Mar/2025 02: 56PM
Lab No : 10140489	Sample Collected ON : 10/Mar/2025 02: 57PM
Referred By : Dr.YUSUF ANSARI	Sample Received ON : 10/Mar/2025 03: 21PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 10/Mar/2025 04: 52PM
Doctor Advice : CPK - MB,TROPONIN-T hs Stat,2D ECHO	



Test Name	Result	Unit	Bio. Ref. Range	Method
TROPONIN-T hs Stat				
TROPONIN-T	0.007	ng/ml	< 0.010	

NOTES :-

Troponin T hs is a member of the myofibrillar proteins of striated muscularis. These myofibrillar proteins are the building blocks of the contractile apparatus. Troponin T hs binds the troponin complex to tropomyosin and binds the neighboring tropomyosin molecules. The determination of troponin T in serum plays an important role in the diagnosis of myocardial infarction(AMI),microinfarction (minor myocardial damage - MMO) and myocarditis.Troponin T is detectable about 3 -4 hours after the occurrence of cardiac symptoms .Following acute myocardial ischemia ,Troponin T remains in the serum for a lengthy period of time and can hence help to detect myocardial events that have occurred upto 14 days earlier.

Cobas E 411 Troponin T hs Stat employs monoclonal antibodies specifically directed against human cardiac Troponin T (after release from the free cytosol and myofibrils .)

Based on the WHO criteria for the definition of AMI from the 1970~s the cutoff (clinical discriminator) value for troponin T is 0.1 ng/ml according to ROC analysis.

Elevated Troponin T values are occasionally found in patients with restricted renal function despite the absence of definite evidence of myocardial Ischemia.

(ELECTRO-CHEMILUMINESCENCE TECHNIQUE BY Cobas E 411)

*** End Of Report ***

CHARAK

[Checked By]

Print.Date/Time: 10-03-2025 18:17:49

*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

DR. ADITI D AGARWAL
PATHOLOGIST

Aditi D Agarwal

Patient Name	: Mr.TAJUDDIN	Visit No	: CHA250043194
Age/Gender	: 55 Y/M	Registration ON	: 10/Mar/2025 02:56PM
Lab No	: 10140489	Sample Collected ON	: 10/Mar/2025 02:56PM
Referred By	: Dr.YUSUF ANSARI	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 10/Mar/2025 05:38PM

2D- ECHO & COLOR DOPPLER REPORT

1. **MITRAL VALVE STUDY** : MVOA - Normal (perimetry) cm² (PHT)

Anterior Mitral Leaflet:

- (a) **Motion**: Normal (b) **Thickness** : Normal (c) **DE** : 189 cm.
 (d) **EF** : 93 mm/sec (e) **EPSS** : 06 mm (f) **Vegetation** : -
 (g) **Calcium** : -

Posterior mitral leaflet : Normal

- (a). **Motion** : Normal (b) **Calcium**: - (c) **Vegetation** : -

Valve Score : Mobility /4 Thickness /4 SVA /4
 Calcium /4 Total /16

2. **AORTIC VALVE STUDY**

- (a) **Aortic root** : 2.8cms (b) **Aortic Opening** : 2.0cms (c) **Closure**: Central
 (d) **Calcium** : - (e) **Eccentricity Index** : 1 (f) **Vegetation** : -

(g) **Valve Structure** : Tricuspid,

3. **PULMONARY VALVE STUDY** Normal

- (a) **EF Slope** : - (b) **A Wave** : + (c) **MSN** : -

(D) **Thickness** : (e) **Others** :

4. **TRICUSPID VALVE** : Normal

5. **SEPTAL AORTIC CONTINUITY** 6. **AORTIC MITRAL CONTINUITY**

Left Atrium : 2.5 cms

Clot : -

Others :

Right Atrium : Normal

Clot : -

Others : -

Contd.....



Patient Name	: Mr.TAJUDDIN	Visit No	: CHA250043194
Age/Gender	: 55 Y/M	Registration ON	: 10/Mar/2025 02:56PM
Lab No	: 10140489	Sample Collected ON	: 10/Mar/2025 02:56PM
Referred By	: Dr.YUSUF ANSARI	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 10/Mar/2025 05:38PM

VENTRICLES

RIGHT VENTRICLE : Normal

RVD (D)

RVOT

LEFT VENTRICLE :

LVIVS (D) 0.8 cm (s)1.6 cm

Motion : normal

LVPW (D) 0.8cm (s) 1.6 cm

Motion : Normal

LVID (D) 4.5 cm (s) 2.4 cm

Ejection Fraction :76%

Fractional Shortening : 45 %

TOMOGRAPHIC VIEWS

Parasternal Long axis view :

NORMAL LV RV DIMENSION
GOOD LV CONTRACTILITY.

Short axis view

Aortic valve level :

AOV - NORMAL
PV - NORMAL
TV - NORMAL

Mitral valve level :

MV - NORMAL

Papillary Muscle Level :

NO RWMA

Apical 4 chamber View :

No LV CLOT



Patient Name	: Mr.TAJUDDIN	Visit No	: CHA250043194
Age/Gender	: 55 Y/M	Registration ON	: 10/Mar/2025 02:56PM
Lab No	: 10140489	Sample Collected ON	: 10/Mar/2025 02:56PM
Referred By	: Dr.YUSUF ANSARI	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 10/Mar/2025 05:38PM

PERICARDIUM

Normal

DOPPLER STUDIES

	Velocity (m/sec)	Flow pattern (/4)	Regurgitation	Gradient (mm Hg)	Valve area (cm 2)
MITRAL	e = 0.8 a = 0.6	Normal	-	-	-
AORTIC	1.4	Normal	-	-	-
TRICUSPID	0.4	Normal	-	-	-
PULMONARY	1.1	Normal	-	-	-

OTHER HAEMODYNAMIC DATA

COLOUR DOPPLER

NO REGURGITATION OR TURBULENCE ACROSS ANY VALVE

CONCLUSIONS :

- NORMAL LV RV DIMENSION
- GOOD LV SYSTOLIC FUNCTION
- LVEF = 76 %
- NO RWMA
- ALL VALVES NORMAL
- NO CLOT / VEGETATION
- NO PERICARDIAL EFFUSION

OPINION – NORMAL 2D-ECHO & COLOUR DOPPLER STUDY

DR. RAJIV RASTOGI, MD,DM

*** End Of Report ***

