

Patient Name	: Mr.RAM JI	Visit No	: CHA250043270
Age/Gender	: 40 Y/M	Registration ON	: 10/Mar/2025 04:44PM
Lab No	: 10140565	Sample Collected ON	: 10/Mar/2025 04:44PM
Referred By	: Dr.KGMU	Sample Received ON	:
Refer Lab/Hosp	: CHARAK NA	Report Generated ON	: 10/Mar/2025 06:33PM

CECT STUDY OF FACE & NECK

- A lobulated heterogeneously enhancing soft tissue attenuation lesion measuring approx. 68 x 22 x 41 mm is seen involving left buccal mucosa with contiguous involvement of left superior & inferior gingivo buccal sulci. Posteriorly it shows extension into left retromolar trigone & masticator space region with infiltration into left masseter, temporalis and medial pterygoid muscles. It is causing permeative erosion of angle of left hemimandible. Further medially it shows extension into left parapharyngeal fat space & pharyngeal mucosal space with infiltration into left tonsillar fossa and tonsillar pillars which appear bulky. Laterally there is extension into left buccal space with encasement of left common fascial vein and infiltration into overlying skin.
- Few variable sized heterogeneously enhancing left cervical lymphnodes are seen at levels Ib & II with the largest measuring approx. 16 mm in MSAD seen at left level Ib.
- Bilateral submandibular and parotid glands are seen normally.
- The aryepiglottic folds and epiglottis are seen normally.
- The valleculae and pyriform sinuses are seen normally.
- The laryngeal airway and cartilages are seen normally.
- Degenerative changes are seen in visualized parts of spine. No evidence of any lytic or sclerotic area is seen.
- Thyroid gland is normal in size and density.

IMPRESSION:

KNOWN CASE OF CARCINOMA LEFT BUCCAL MUCOSA SHOWING : -

- HETEROGENEOUSLY ENHANCING LESION INVOLVING LEFT BUCCAL MUCOSA COMPLEX WITH FEATURES, EXTENSIONS, OSTEO-EROSION AND CERVICAL LYMPHADENOPATHY AS DESCRIBED.

Clinical correlation is necessary.

(DR. JAYENDRA KUMAR, MD)

Transcribed by Rachna

*** End Of Report ***

