

292/05, Tulsidas Marg, Basement Chowk, Lucknow-226 003

Phone: 0522-4062223, 9305548277, 8400888844 9415577933, 9336154100, Tollfree No.: 8688360360

E-mail: charak1984@gmail.com

CMO Reg. No. RMEE 2445133 NABL Reg. No. MC-2491 Certificate No. MIS-2023-0218

Patient Name : Ms.FATIMA Visit No : CHA250043898

 Age/Gender
 : 65 Y/F
 Registration ON
 : 11/Mar/2025 01:32PM

 Lab No
 : 10141193
 Sample Collected ON
 : 11/Mar/2025 01:32PM

Referred By : Dr.AFTAB ALAM KHAN Sample Received ON :

Refer Lab/Hosp : CHARAK NA Report Generated ON : 11/Mar/2025 03:18PM

ULTRASOUND STUDY OF WHOLE ABDOMEN

Excessive gaseous abdomen (poor bowel preparation)

- <u>Liver</u> is mildly enlarged in size (~150mm) and shows homogenous echotexture of liver parenchyma. No intrahepatic biliary radicle dilatation is seen. No space occupying lesion is seen. Hepatic veins and IVC are seen normally.
- Gall bladder is normal in size and shows few calculi in GB lumen measuring upto approx 1.9mm. No mass lesion is seen. GB wall is thickened measuring approx 3.1mm.
- CBD is normal at porta. No obstructive lesion is seen.
- Portal vein Portal vein is normal at porta.
- <u>Pancreas</u> is normal in size and shows homogenous echotexture of parenchyma. PD is not dilated. No parenchymal calcification is seen. No peripancreatic collection is seen.
- <u>Spleen</u> is normal in size and shows homogenous echotexture of parenchyma. No SOL is seen.
- No retroperitoneal adenopathy is seen.
- <u>Both kidneys</u> are normal in size and position. No hydronephrosis is seen. No calculus or mass lesion is seen. Cortico-medullary differentiation is well maintained. Parenchymal thickness is normal. No scarring is seen. Right kidney measures 83 x 35 mm in size. Left kidney measures 86 x 37 mm in size.
- <u>Ureters</u> Both ureters are not dilated. UVJ are seen normally.
- <u>Urinary bladder</u> is not distended. Foley's catheter is seen in situ.
- Bilateral minimal pleural effusion is seen.
- Mildly dilated bowel loops, filled with semisolid content with hypokinetic bowel movement with minimal inter bowel fluid collection in peritoneal cavity.

OPINION:

PR

CHARAK

- MILD HEPATOMEGALY.
- CHOLELITHIASIS WITH CHOLECYSTITIS.
- BILATERAL MINIMAL PLEURAL EFFUSION.
- MILDLY DILATED BOWEL LOOPS, FILLED WITH SEMISOLID CONTENT WITH HYPOKINETIC BOWEL MOVEMENT WITH MINIMAL INTER BOWEL FLUID COLLECTION IN PERITONEAL CAVITY S.A.I.O (ADV: X- RAY ERECT ABDOMEN TO R/O S.A.I.O).

Clinical correlation is necessary.

[DR. R. K. SINGH, MD]

Transcribed by Gausiya





292/05, Tulsidas Marg, Basement Chowk, Lucknow-226 003 **Phone**: 0522-4062223, 9305548277, 8400888844

9415577933, 9336154100, **Tollfree No.**: 8688360360 **E-mail** : charak1984@gmail.com

CMO Reg. No. RMEE 2445133 NABL Reg. No. MC-2491 Certificate No. MIS-2023-0218

Patient Name : Ms.FATIMA Visit No : CHA250043898

 Age/Gender
 : 65 Y/F
 Registration ON
 : 11/Mar/2025 01:32PM

 Lab No
 : 10141193
 Sample Collected ON
 : 11/Mar/2025 01:32PM

Referred By : Dr.AFTAB ALAM KHAN Sample Received ON

Refer Lab/Hosp : CHARAK NA Report Generated ON : 11/Mar/2025 02:52PM

SKIAGRAM ABDOMEN (ERECT) AP VIEW

• No free gas is seen under both dome of diaphragm.

- No abnormal air fluid levels are seen.
- Ryle's tube is seen in situ .
- Contrast filled large bowel loops are seen [post CT] .

Clinical correlation is necessary.

[DR. RAJESH KUMAR SHARMA, MD]

TRANSCRIBED BY: ANUP

*** End Of Report ***



