

Patient Name : Mr. JK VERMA	Visit No : CHA250045242
Age/Gender : 68 Y/M	Registration ON : 13/Mar/2025 01: 17PM
Lab No : 10142537	Sample Collected ON : 13/Mar/2025 01: 19PM
Referred By : Dr. UNIQUE HOSPITAL	Sample Received ON : 13/Mar/2025 01: 19PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 13/Mar/2025 02: 43PM
Doctor Advice : USG WHOLE ABDOMEN, URINE COM. EXMAMINATION, PSA-TOTAL, MCU, RGU	



Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

URINE EXAMINATION REPORT

Colour-U	STRAW		Light Yellow	
Appearance (Urine)	CLEAR		Clear	
Specific Gravity	1.010		1.005 - 1.025	
pH-Urine	Acidic (6.5)		4.5 - 8.0	
PROTEIN	10 mg/dl	mg/dl	ABSENT	Dipstick
Glucose	Absent		Absent	
Ketones	Absent		Absent	
Bilirubin-U	Absent		Absent	
Blood-U	PRESENT		Absent	
Urobilinogen-U	0.20	EU/dL	0.2 - 1.0	
Leukocytes-U	Absent		Absent	
NITRITE	Absent		Absent	
MICROSCOPIC EXAMINATION				
Pus cells / hpf	Nil	/hpf	< 5/hpf	
Epithelial Cells	1-2	/hpf	0 - 5	
RBC / hpf	4-6		< 3/hpf	

CHARAK

[Checked By]

Print.Date/Time: 13-03-2025 20:00:10

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)

Patient Name : Mr. JK VERMA	Visit No : CHA250045242
Age/Gender : 68 Y/M	Registration ON : 13/Mar/2025 01:17PM
Lab No : 10142537	Sample Collected ON : 13/Mar/2025 01:19PM
Referred By : Dr. UNIQUE HOSPITAL	Sample Received ON : 13/Mar/2025 01:52PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 13/Mar/2025 04:11PM
Doctor Advice : USG WHOLE ABDOMEN, URINE COM. EXMAMINATION, PSA-TOTAL, MCU, RGU	



Test Name	Result	Unit	Bio. Ref. Range	Method
PSA-TOTAL				
PROSTATE SPECIFIC ANTIGEN	67.3	ng/mL	0.2-4.0	CLIA

NOTE – Findings checked twice. Please correlate clinically.

COMMENT : 1. Prostate specific antigen (PSA) is useful for diagnosis of disseminated CA prostate & its sequential measurement is the most sensitive measure of monitoring treatment of disseminated CA prostate with its shorter half life (half life of 2.2 days only) it is superior to prostatic acid phosphatase(PAP). PSA is elevated in nearly all patients with stage D carcinoma whereas PAP is elevated in only 45 % of patient. Mild PSA elevation are also reported in some patients of BHP.
2. Blood samples should be obtained before prostate biopsy or prostatectomy or prostatic massage or digital pre rectal examination as it may result intrasient levation of PSA value for few days.

NOTE :- PSA values obtained in different types of PSA assay methods cannot be used interchangeably as the PSA value in a given sample varies with assays from different manufactures due to difference in assay methodology and reagent specificity. If in the course of monitoring a patient the assay method used for determination is changed, additional sequential testing should be carried out to confirm baseline value.

DONE BY;
Enhanced Chemiluminescence "VITROS ECI"

*** End Of Report ***



[Checked By]

