

Patient Name : Mr.EHTISHAMUL HAQ	Visit No : CHA250046491
Age/Gender : 66 Y/M	Registration ON : 16/Mar/2025 04: 26PM
<b>Lab No : 10143786</b>	Sample Collected ON : 16/Mar/2025 04: 27PM
Referred By : Dr.ROHAN BAJPAI	Sample Received ON : 16/Mar/2025 04: 27PM
Refer Lab/Hosp : CGHS (BILLING)	Report Generated ON : 16/Mar/2025 05: 47PM
Doctor Advice : PSA-TOTAL,CALCIUM,URINE COM. EXMAMINATION,KIDNEY FUNCTION TEST - I,CBC+ESR	



Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

**URINE EXAMINATION REPORT**

Colour-U	<b>Light yellow</b>		Light Yellow	
Appearance (Urine)	CLEAR		Clear	
Specific Gravity	<b>1.015</b>		1.005 - 1.025	
pH-Urine	Acidic (6.0)		4.5 - 8.0	
PROTEIN	20 mg/dl	mg/dl	ABSENT	Dipstick
Glucose	Absent		Absent	
Ketones	Absent		Absent	
Bilirubin-U	Absent		Absent	
Blood-U	Absent		Absent	
Urobilinogen-U	0.20	EU/dL	0.2 - 1.0	
Leukocytes-U	Absent		Absent	
NITRITE	Absent		Absent	
<b>MICROSCOPIC EXAMINATION</b>				
Pus cells / hpf	Occasional	/hpf	< 5/hpf	
Epithelial Cells	2-4	/hpf	0 - 5	
RBC / hpf	Nil		< 3/hpf	

CHARAK

[Checked By]

Print.Date/Time: 17-03-2025 00:00:47

\*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA  
PATHOLOGIST

DR. SHADAB  
PATHOLOGIST

*Aditi D Agarwal*  
DR. ADITI D AGARWAL  
PATHOLOGIST

Patient Name : Mr.EHTISHAMUL HAQ	Visit No : CHA250046491
Age/Gender : 66 Y/M	Registration ON : 16/Mar/2025 04: 26PM
<b>Lab No : 10143786</b>	Sample Collected ON : 16/Mar/2025 04: 27PM
Referred By : Dr.ROHAN BAJPAI	Sample Received ON : 16/Mar/2025 04: 29PM
Refer Lab/Hosp : CGHS (BILLING)	Report Generated ON : 16/Mar/2025 05: 28PM
Doctor Advice : PSA-TOTAL,CALCIUM,URINE COM. EXMAMINATION,KIDNEY FUNCTION TEST - I,CBC+ESR	



Test Name	Result	Unit	Bio. Ref. Range	Method
<b>PSA-TOTAL</b>				
PROSTATE SPECIFIC ANTIGEN	<b>0.15</b>	ng/mL	0.2-4.0	CLIA

COMMENT : 1. Prostate specific antigen (PSA) is useful for diagnosis of disseminated CA prostate & its equential measurement is the most sensitive measure of monitoring treatment of disseminated CA prostate with its shorter half life (half life of 2.2 days only) it is superior to prostatic acis phosphatase(PAP). PSA is elevated in nearly all patients with stage D carcinoma whereas PAP is elevated in only 45 % of patient. Mild PSA elevation are also reported in some patients of BHP.

2. Blood samples should be obtained before prostate biopsy or prostatectomy or prostatic massage or digital pre rectal examination as it may result intrasient levation of PSA value for few days.

NOTE :- PSA values obtained in different types of PSA assay methods cannot be used interchangeably as the PSA value in a given sample varies with assays from different manufactures due to difference in assay methodology and reagent specificity. If in the course of monitoring a patient the assay method used for determination is changed, additional sequential testing should be carried out to confirm baseline value.

DONE BY;  
Enhanced Chemiluminescence "VITROS ECI"

\*\*\* End Of Report \*\*\*

CHARAK



[Checked By]



DR. NISHANT SHARMA  
PATHOLOGIST

DR. SHADAB  
PATHOLOGIST

*Dr. Aditi D Agarwal*  
DR. ADITI D AGARWAL  
PATHOLOGIST