


Patient Name : Mr.DAYAL SHARAN SINGH	Visit No : CHA250046927
Age/Gender : 75 Y/M	Registration ON : 17/Mar/2025 11:33AM
Lab No : 10144222	Sample Collected ON : 17/Mar/2025 11:50AM
Referred By : Dr.WE CARE HOSPITAL	Sample Received ON : 17/Mar/2025 12:07PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 04:28PM
Doctor Advice : 2D ECHO,USG KUB,PSA-TOTAL,URIC ACID,UACR	



Test Name	Result	Unit	Bio. Ref. Range	Method
URIC ACID				
Sample Type : SERUM				
SERUM URIC ACID	8.4	mg/dL	2.40 - 5.70	Uricase,Colorimetric
URINE ALBUMIN CREATININE RATIO				
URINE FOR MICRO ALBUMIN	24	MG/L	< 20 MG/L	
URINARY CREATININE	86.80	mg/dL	20-320 mg/dL	
URINE ALBUMIN CREATININE RATIO	27.6	mg/g		calculated



CHARAK



[Checked By]

Print.Date/Time: 17-03-2025 17:11:46

*Patient Identity Has Not Been Verified. Not For Medicolegal

DR. NISHANT SHARMA PATHOLOGIST
DR. SHADAB PATHOLOGIST
DR. ADITI D AGARWAL PATHOLOGIST

Patient Name : Mr.DAYAL SHARAN SINGH	Visit No : CHA250046927
Age/Gender : 75 Y/M	Registration ON : 17/Mar/2025 11:33AM
Lab No : 10144222	Sample Collected ON : 17/Mar/2025 11:50AM
Referred By : Dr.WE CARE HOSPITAL	Sample Received ON : 17/Mar/2025 12:07PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 02:31PM
Doctor Advice : 2D ECHO,USG KUB,PSA-TOTAL,URIC ACID,UACR	



Test Name	Result	Unit	Bio. Ref. Range	Method
PSA-TOTAL				
PROSTATE SPECIFIC ANTIGEN	0.4	ng/mL	0.2-4.0	CLIA

COMMENT : 1. Prostate specific antigen (PSA) is useful for diagnosis of disseminated CA prostate & its equential measurement is the most sensitive measure of monitoring treatment of disseminated CA prostate with its shorter half life (half life of 2.2 days only) it is superior to prostatic acis phosphatase(PAP). PSA is elevated in nearly all patients with stage D carcinoma whereas PAP is elevated in only 45 % of patient. Mild PSA elevation are also reported in some patients of BHP.

2. Blood samples should be obtained before prostate biopsy or prostatecomy or prostatic massage or digital pre rectal examination as it may result intrasient levation of PSA value for few days.

NOTE :- PSA values obtained in different types of PSA assay methods cannot be used interchangeably as the PSA value in a given sample varies with assays from different manufactures due to difference in assay methodology and reagent specificity. If in the course of monitoring a patient the assay method used for determination is changed, additional sequential testing should be carried out to confirm baseline value.

DONE BY;
Enhanced Chemiluminescence "VITROS ECI"

*** End Of Report ***

CHARAK



MC-2491

Print.Date/Time: 17-03-2025 17:11:48

*Patient Identity Has Not Been Verified. Not For Medicolegal



Sharma

DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

Dr. SYED SAIF AHMAD
MD (MICROBIOLOGY)