

Patient Name : Mr.MOHD HASIM	Visit No : CHA250047100
Age/Gender : 35 Y/M	Registration ON : 17/Mar/2025 01:07PM
Lab No : 10144395	Sample Collected ON : 17/Mar/2025 01:11PM
Referred By : Dr.MANISH TANDON	Sample Received ON : 17/Mar/2025 01:23PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 04:23PM
Doctor Advice : USG WHOLE ABDOMEN,HBSAg,HCV,HIV,T3T4TSH,FASTING,NA+K+,CREATININE,Albumin,LFT,CRP (Quantitative),ESR,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
ESR				
Erythrocyte Sedimentation Rate ESR	8.00		0 - 15	Westergreen

Note:

1. Test conducted on EDTA whole blood at 37°C.
2. ESR readings are auto- corrected with respect to Hematocrit (PCV) values.
3. It indicates presence and intensity of an inflammatory process. It is a prognostic test and used to monitor the course or response to treatment of diseases like tuberculosis, acute rheumatic fever. It is also increased in multiple myeloma, hypothyroidism.

CRP-QUANTITATIVE

CRP-QUANTITATIVE TEST	4.3	MG/L	0.1 - 6
-----------------------	-----	------	---------

Method: Immunoturbidimetric

(Method: Immunoturbidimetric on photometry system)

SUMMARY : C - reactive protien (CRP) is the best known among the acute phase protiens, a group of protien whose concentration increases in blood as a response to inflammatory disorders.CRP is normally present in low concentration in blood of healthy individuals (< 1mg/L). It is elevated up to 500 mg/L in acute inflammatory processes associated with bacterial infections, post operative conditions tissue damage already after 6 hours reaching a peak at 48 hours.. The measurement of CRP represents a useful laboratory test for detection of acute infection as well as for monitoring inflammtory proceses also in acute rheumatic & gastrointestinal disease. In recent studies it has been shows that in apparrently healthy subjects there is a direct orrelation between CRP concentrations & the risk of developing oronary heart disease (CHD).

hsCRP cut off for risk assessment as per CDC/AHA

Level	Risk
<1.0	Low
1.0-3.0	Average
>3.0	High

All reports to be clinically corelated

SERUM ALBUMIN

ALBUMIN	4.8	gm/dl	3.20 - 5.50	Bromcresol Green (BCG)
---------	-----	-------	-------------	------------------------



[Checked By]

Print.Date/Time: 17-03-2025 17:26:24

*Patient Identity Has Not Been Verified. Not For Medicolegal

DR. NISHANT SHARMA PATHOLOGIST
DR. SHADAB PATHOLOGIST
DR. ADITI D AGARWAL PATHOLOGIST

Patient Name : Mr.MOHD HASIM	Visit No : CHA250047100
Age/Gender : 35 Y/M	Registration ON : 17/Mar/2025 01:07PM
Lab No : 10144395	Sample Collected ON : 17/Mar/2025 01:11PM
Referred By : Dr.MANISH TANDON	Sample Received ON : 17/Mar/2025 01:23PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 04:23PM
Doctor Advice : USG WHOLE ABDOMEN,HBSAg,HCV,HIV,T3T4TSH,FASTING,NA+K+,CREATININE,Albumin,LFT,CRP (Quantitative),ESR,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
HEPATITIS B SURFACE ANTIGEN (HBsAg)				
Sample Type : SERUM				
HEPATITIS B SURFACE ANTIGEN	NON REACTIVE		<1 - Non Reactive >1 - Reactive	CMIA

Note: This is only a Screening test. Confirmation of the result (Non Reactive/Reactive)should be done by performing a PCR based test.

COMMENTS:

-HBsAg is the first serological marker after infection with Hepatitis B Virus appearing one to ten weeks after exposure and two to eight weeks before the onset of clinical symptoms. HBsAg persists during the acute phase and clears late in the convalescence phase. Failure to clear HBsAg within six months indicates a chronic HBsAg carrier state. HBsAg assays are used to identify the persons infected with HBV and to prevent transmission of the virus by blood and blood products as well as to monitor the status of infected individuals in combination with other hepatitis B serological markers.
-Borderline cases must be confirmed with confirmatory neutralizing assay.

LIMITATIONS:

-Results should be used in conjunction with patient history and other hepatitis markers for diagnosis of acute and chronic infections.
-Specimens from patients who have received preparations of mouse monoclonal antibodies for diagnosis or therapy may contain human anti-mouse antibodies (HAMA) which may produce anomalous values when tested with assay kits that employs mouse monoclonal antibodies.
-Heterophilic antibodies in human serum can react with reagent immunoglobulins, interfering with in vitro immunoassays. Patients routinely exposed to animals or animal serum products can be prone to this interference and anomalous results may be observed.
-Cross reactivity for specimens from individual with medical conditions (Pregnancy, HIV etc) has been observed.
-HBsAg mutations may result in a false negative result in some HBsAg assays.
-If HBsAg results are inconsistent with clinical evidence, additional testing is suggested to confirm the result.



HIV				
HIV-SEROLOGY	NON REACTIVE		<1.0 : NON REACTIVE >1.0 : REACTIVE	

Done by: Vitros ECI (Sandwich Assay)

Note:-Elisa test is a screening method for HIV.It is known to give false Positive & Negative result.
Hence confirmation:"Western Blot" method is advised.



[Checked By]

Print.Date/Time: 17-03-2025 17:26:25

*Patient Identity Has Not Been Verified. Not For Medicolegal

DR. NISHANT SHARMA PATHOLOGIST
DR. SHADAB PATHOLOGIST
DR. ADITI D AGARWAL PATHOLOGIST


Patient Name : Mr.MOHD HASIM	Visit No : CHA250047100
Age/Gender : 35 Y/M	Registration ON : 17/Mar/2025 01:07PM
Lab No : 10144395	Sample Collected ON : 17/Mar/2025 01:11PM
Referred By : Dr.MANISH TANDON	Sample Received ON : 17/Mar/2025 01:23PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 04:23PM
Doctor Advice : USG WHOLE ABDOMEN,HBSAg,HCV,HIV,T3T4TSH,FASTING,NA+K+,CREATININE,Albumin,LFT,CRP (Quantitative),ESR,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
HCV				
Anti-Hepatitis C Virus Antibodies.	NON REACTIVE		< 1.0 : NON REACTIVE > 1.0 : REACTIVE	Sandwich Assay

Done by: Vitros ECI (Sandwich Assay)

Note:This is only a Screening test. Confirmation of the result (Non Reactive/Reactive)should be done by performing a PCR based test.



CHARAK

[Checked By]

Print.Date/Time: 17-03-2025 17:26:26

*Patient Identity Has Not Been Verified. Not For Medicolegal



DR. NISHANT SHARMA
PATHOLOGIST

DR. SHADAB
PATHOLOGIST

Dr. Aditi D Agarwal
DR. ADITI D AGARWAL
PATHOLOGIST

Patient Name : Mr.MOHD HASIM	Visit No : CHA250047100
Age/Gender : 35 Y/M	Registration ON : 17/Mar/2025 01:07PM
Lab No : 10144395	Sample Collected ON : 17/Mar/2025 01:11PM
Referred By : Dr.MANISH TANDON	Sample Received ON : 17/Mar/2025 01:51PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 03:49PM
Doctor Advice : USG WHOLE ABDOMEN,HBSAg,HCV,HIV,T3T4TSH,FASTING,NA+K+,CREATININE,Albumin,LFT,CRP (Quantitative),ESR,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
CBC (COMPLETE BLOOD COUNT)				
Hb	15.5	g/dl	12 - 15	Non Cyanide
R.B.C. COUNT	5.50	mil/cmm	3.8 - 4.8	Electrical Impedence
PCV	48.2	%	36 - 45	Pulse hieght detection
MCV	87.0	fL	80 - 96	calculated
MCH	28.0	pg	27 - 33	Calculated
MCHC	32.2	g/dL	30 - 36	Calculated
RDW	13.2	%	11 - 15	RBC histogram derivation
RETIC	0.4 %	%	0.5 - 2.5	Microscopy
TOTAL LEUCOCYTES COUNT	9840	/cmm	4000 - 10000	Flocytometry
DIFFERENTIAL LEUCOCYTE COUNT				
NEUTROPHIL	62	%	40 - 75	Flowcytometry
LYMPHOCYTES	33	%	25 - 45	Flowcytometry
EOSINOPHIL	2	%	1 - 6	Flowcytometry
MONOCYTE	3	%	2 - 10	Flowcytometry
BASOPHIL	0	%	00 - 01	Flowcytometry
PLATELET COUNT	360,000	/cmm	150000 - 450000	Elect Imped..
PLATELET COUNT (MANUAL)	360000	/cmm	150000 - 450000	Microscopy .
Absolute Neutrophils Count	6,101	/cmm	2000 - 7000	Calculated
Absolute Lymphocytes Count	3,247	/cmm	1000-3000	Calculated
Absolute Eosinophils Count	197	/cmm	20-500	Calculated
Absolute Monocytes Count	295	/cmm	200-1000	Calculated
Mentzer Index	16			
Peripheral Blood Picture	:			

Red blood cells are normocytic normochromic. Platelets are adequate. No immature cells or parasite seen.



[Checked By]



Patient Name : Mr.MOHD HASIM	Visit No : CHA250047100
Age/Gender : 35 Y/M	Registration ON : 17/Mar/2025 01:07PM
Lab No : 10144395	Sample Collected ON : 17/Mar/2025 01:11PM
Referred By : Dr.MANISH TANDON	Sample Received ON : 17/Mar/2025 01:23PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 03:11PM
Doctor Advice : USG WHOLE ABDOMEN,HBSAg,HCV,HIV,T3T4TSH,FASTING,NA+K+,CREATININE,Albumin,LFT,CRP (Quantitative),ESR,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
FASTING				
Blood Sugar Fasting	170.8	mg/dl	70 - 110	Hexokinase
NA+K+				
SODIUM Serum	136.0	MEq/L	135 - 155	ISE Direct
POTASSIUM Serum	4.2	MEq/L	3.5 - 5.5	ISE Direct
SERUM CREATININE				
CREATININE	0.90	mg/dl	0.50 - 1.40	Alkaline picrate-kinetic
LIVER FUNCTION TEST				
TOTAL BILIRUBIN	0.78	mg/dl	0.4 - 1.1	Diazonium Ion
CONJUGATED (D. Bilirubin)	0.10	mg/dL	0.00-0.30	Diazotization
UNCONJUGATED (I.D. Bilirubin)	0.68	mg/dL	0.1 - 1.0	Calculated
ALK PHOS	99.60	U/L	30 - 120	PNPP, AMP Buffer
SGPT	23.0	U/L	5 - 40	UV without P5P
SGOT	17.0	U/L	5 - 40	UV without P5P

CHARAK



[Checked By]



Sharma

DR. NISHANT SHARMA PATHOLOGIST DR. SHADAB PATHOLOGIST Dr. SYED SAIF AHMAD MD (MICROBIOLOGY)

Patient Name : Mr.MOHD HASIM	Visit No : CHA250047100
Age/Gender : 35 Y/M	Registration ON : 17/Mar/2025 01:07PM
Lab No : 10144395	Sample Collected ON : 17/Mar/2025 01:11PM
Referred By : Dr.MANISH TANDON	Sample Received ON : 17/Mar/2025 01:23PM
Refer Lab/Hosp : CHARAK NA	Report Generated ON : 17/Mar/2025 02:36PM
Doctor Advice : USG WHOLE ABDOMEN,HBSAg,HCV,HIV,T3T4TSH,FASTING,NA+K+,CREATININE,Albumin,LFT,CRP (Quantitative),ESR,CBC (WHOLE BLOOD)	



Test Name	Result	Unit	Bio. Ref. Range	Method
T3T4TSH				
T3	2.35	nmol/L	1.49-2.96	ECLIA
T4	120.76	n mol/l	63 - 177	ECLIA
TSH	3.55	uIU/ml	0.47 - 4.52	ECLIA

Note

- (1) Patients having low T3 & T4 levels but high TSH levels suffer from primary hypothyroidism,cretinism,juvenile mysedema or autoimmune disorders.
- (2) Patients having low T3 & T4 levels but high TSH levels suffer from grave~s disease, toxic adenoma or sub-acute thyroiditis.
- (3) Patients having either low or normal T3 & T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.
- (4) Patients having high T3 & T4 levels but normal TSH levels may suffer from toxic multinodular goitre. This condition is mostly asymptomatic and may cause transient hyperthyroidism but no persistent symptoms.
- (5) Patient with high or normal T3 & T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 Toxicosis respectively.
- (6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the cacabolic state and may revert tonormal when the patient recovers.
- (7) There are many drugs for eg.Glucocorticoids ,dopamine,Lithium,iodides ,oral radiographic dyes,ets.Which may affect the thyroid function tests.
- (8) Generally when total T3& T4 results are indecisive then Free T3 & Free T4 test are recommended for further confirmation along with

(1 Beckman Dxi-600 2. ELECTRO-CHEMILUMINESCENCE TECHINIQUE BY ELECSYSYS -E411)

*** End Of Report ***

CHARAK



[Checked By]



Sham

DR. NISHANT SHARMA DR. SHADAB Dr. SYED SAIF AHMAD
PATHOLOGIST PATHOLOGIST MD (MICROBIOLOGY)