



itDose

Taking You Beyond The Limits

Visit ID : ITD1369	Registration : 06/Feb/2023 04: 20PM
UHID/MR No : ITD.0000001231	Collected : 06/Feb/2023 04: 21PM
Patient Name : MR. PASCHIM	Received : 06/Feb/2023 04: 21PM
Age/Gender : 34 Y O M O D /M	Reported : 06/Feb/2023 04: 22PM
Ref Doctor : SELF	Status : Final Report
Client Name : STANDARD	Client Code : 78
Client Add : D 159, 1st Floor, Sector 7, No	Barcode No : 10060011

DEPARTMENT OF BIOCHEMISTRY

SERUM LIPASE					
Sample Type : Serum					
SERUM LIPASE	34	L	U/L	<=38 U/L	Color
COMMENTS: as, such as acute pancreatitis, chronic pancreatitis, and obstruction of the pancreatic duct. In acute pancreatitis serum lipase activity tends to become elevated & remains for about 7 - 10 days .Increased lipase activity rarely lasts longer than 14 days, and prolonged increases suggest a poor prognosis or the presence of a cyst. Serum lipase may also be elevated in patients with chronic pancreatitis, obstruction of the pancreatic duct and non pancreatic conditions including renal diseases, various abdominal diseases such as acute cholecystitis, intestinal obstruction or infarction, duodenal ulcer, and liver disease, as well as alcoholism & diabetic keto-acidosis & in patients who have undergone endoscopic r					
Lipase measurements are used in the diagnosis and treatment of diseases of the pancre					
etrograde cholangiopancreatography. Elevation of serum lipase activity in patients with mumps strongly suggests significant pancreatic as well as salivary gland involvement by the disease.....					

*** End Of Report ***

Prhose

Dr. Pritha Ghose
MBBS, MD (Pathology)
Consultant Pathologist

Dr. Sourav Gangopadhyay
MBBS, MD (Pathology)
Consultant Pathologist

Dr. Ranjan Basu
MBBS, MD (Microbiology)
Consultant Microbiologist

Dr. Shrinwanti Karmakar
MBBS, MD (Biochemistry)
Consultant Biochemist

