

Visit ID : ITS1440

UHID/MR No : ITD.0000000615

Patient Name : MR. SHIVANGI SHUKLA

Age/Gender : 25 Y 3 M 3 D /F

Client Name

Ref Doctor : SELF

: STANDARD Client Add : D 159, 1st Floor, Sector 7, No Registration : 23/Feb/2023 01:53PM

Collected : 23/Feb/2023 01:54PM

Received : 23/Feb/2023 01:55PM Reported : 23/Feb/2023 04:10PM

Status : Final Report

Client Code : 78 Barcode No : 10060078

DEPARTMENT OF BIOCHEMISTRY

SERUM LIPASE						
Sample Type : Serum						
SERUM LIPASE	5	L	U/L	<=38 U/L	Color	

normal

COMMENTS:as, such as acute pancreatitis, chronic pancreatitis, and obstruction of the pancreatic duct. In acute pancreatitis serum lipase activity tends to become elevated & remains for about 7 - 10 days. Increased lipase activity rarely lasts longer than 14 days, and prolonged increases suggest a poor prognosis or the presence of a cyst. Serum lipase may also be elevated in patients with chronic pancreatitis, obstruction of the pancreatic duct and non pancreatic conditions including renal diseases, various abdominal diseases such as acute cholecystitis, intestinal obstruction or infarction, duodenal ulcer, and liver disease, as well as alcoholism & diabetic keto-acidosis & in patients who have undergone endoscopic r

Lipase measurements are used in the diagnosis and treatment of diseases of the pancre

etrograde cholangiopancreatography. Elevation of serum lipase activity in patients with mumps strongly suggests significant pancreatic as well as salivary gland involvement by the disease......

*** End Of Report ***

Dr. Pritha Ghose MBBS, MD (Pathology)

Consulatant Pathologist

Dr. Sourav Gangopadhyay MBBS, MD (Pathology) Consulatant Pathologist

Dr. Ranjan Basu MBBS, MD (Microbiology) Consultant Microbiologist

Dr. Shrinwanti Karmakar MBBS, MD (Biochemistry) Consultant Biochemist

