



itDose

Taking You Beyond The Limits

Visit ID : ITS1441	Registration : 23/Feb/2023 03: 30PM
UHID/MR No : ITS.0000000017	Collected : 23/Feb/2023 03: 33PM
Patient Name : MR. VIKASH	Received : 23/Feb/2023 03: 35PM
Age/Gender : 35 Y O M O D /M	Reported : 23/Feb/2023 04: 15PM
Ref Doctor : SELF	Status : Final Report
Client Name : STANDARD	Client Code : 78
Client Add : D 159, 1st Floor, Sector 7, No	Barcode No : 10060079

DEPARTMENT OF BIOCHEMISTRY

SERUM LIPASE

Sample Type : Serum

SERUM LIPASE	12	L	U/L	<=38 U/L	Color
--------------	----	---	-----	----------	-------

**COMMENTS:**as, such as acute pancreatitis, chronic pancreatitis, and obstruction of the pancreatic duct. In acute pancreatitis serum lipase activity tends to become elevated & remains for about 7 - 10 days .Increased lipase activity rarely lasts longer than 14 days, and prolonged increases suggest a poor prognosis or the presence of a cyst. Serum lipase may also be elevated in patients with chronic pancreatitis, obstruction of the pancreatic duct and non pancreatic conditions including renal diseases, various abdominal diseases such as acute cholecystitis, intestinal obstruction or infarction, duodenal ulcer, and liver disease, as well as alcoholism & diabetic keto-acidosis & in patients who have undergone endoscopic r

Lipase measurements are used in the diagnosis and treatment of diseases of the pancre

etrograde cholangiopancreatography. Elevation of serum lipase activity in patients with mumps strongly suggests significant pancreatic as well as salivary gland involvement by the disease.....

\*\*\* End Of Report \*\*\*

Dr. Pritha Ghose  
MBBS, MD (Pathology)  
Consultant Pathologist

Dr. Sourav Gangopadhyay  
MBBS, MD (Pathology)  
Consultant Pathologist

Dr. Ranjan Basu  
MBBS, MD (Microbiology)  
Consultant Microbiologist

Dr. Shrinwanti Karmakar  
MBBS, MD (Biochemistry)  
Consultant Biochemist

