

Visit ID : ITS1544

UHID/MR No : ITS.0000000099

Patient Name : TRANS. TTT
Age/Gender : 34 Y 0 M 0 D /T

Ref Doctor

: SELF

Client Name : STANDARD

Client Add : D 159, 1st Floor, Sector 7, No

Registration : 2

Reported

: 25/Mar/2023 11:30AM

Collected : 25/Mar/2023 11:33AM Received : 25/Mar/2023 11:33AM

: 25/Mar/2023 11:33AM : 25/Mar/2023 11:34AM

Status : Final Report

Client Code : 78

Barcode No : 10060175

DEPARTMENT OF BIOCHEMISTRY								
Test Name	Result	Status	Unit	Bio. Ref. Range	Method			

SERUM LIPASE									
Sample Type : Serum									
SERUM LIPASE	2	L	U/L	<=38 U/L	Color				

COMMENTS:as, such as acute pancreatitis, chronic pancreatitis, and obstruction of the pancreatic duct. In acute pancreatitis serum lipase activity tends to become elevated & remains for about 7 - 10 days. Increased lipase activity rarely lasts longer than 14 days, and prolonged increases suggest a poor prognosis or the presence of a cyst. Serum lipase may also be elevated in patients with chronic pancreatitis, obstruction of the pancreatic duct and non pancreatic conditions including renal diseases, various abdominal diseases such as acute cholecystitis, intestinal obstruction or infarction, duodenal ulcer, and liver disease, as well as alcoholism & diabetic keto-acidosis & in patients who have undergone endoscopic r

Lipase measurements are used in the diagnosis and treatment of diseases of the pancre

etrograde cholangiopancreatography. Elevation of serum lipase activity in patients with mumps strongly suggests significant pancreatic as well as salivary gland involvement by the disease......

*** End Of Report ***

Dr. Pritha Ghose MBBS, MD (Pathology) Consulatant Pathologist

Dr. Sourav Gangopadhyay MBBS, MD (Pathology) Consulatant Pathologist Dr. Ranjan Basu MBBS, MD (Microbiology) Consultant Microbiologist Dr. Shrinwanti Karmakar MBBS, MD (Biochemistry) Consultant Biochemist

